

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 850 924 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
14.03.2001 Bulletin 2001/11

(51) Int Cl.7: **C07C 391/02, C07D 333/30,
C07D 233/66, A61K 31/095**

(21) Application number: 97204070.3

(22) Date of filing: 23.12.1997

(54) **Aromatic diselenides and selenosulphides for use as antioxidant agents**

Aromatische Diselenide und Selenosulfide zur Verwendung als Antioxidans-Mittel

Disélenides et selenosulfures utilisables comme agents antioxydant

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

(30) Priority: 27.12.1996 FR 9616103

(43) Date of publication of application:
01.07.1998 Bulletin 1998/27

(73) Proprietor: **OXIS Isle of Man, Limited**
Douglas IM1 3DB, Isle of Man (GB)

(72) Inventors:
• Tailhan-Lomont, Catherine
77310 Boissise-Le-Roi (FR)
• Erdelmeier, Irène
75013 Paris (FR)
• Moutet, Marc
92220 Bagneux (FR)

• Chaudière, Jean
94100 Saint Maur Des Fosses (FR)
• Yadan, Jean-Claude
93100 Montreuil (FR)

(74) Representative: Portal, Gérard et al
Cabinet Beau de Loménie
158, rue de l'Université
75340 Paris Cédex 07 (FR)

(56) References cited:
EP-A- 0 165 534 WO-A-95/27706
US-A- 5 128 365

• P.V. JACQUEMIN, ET AL.: "Synthesis of
2H-3,4-dihydro-1,2-benzoselenazin-3-one and
derivatives: a new heterocyclic ring system"
TETRAHEDRON LETTERS, vol. 33, no. 27, 30
June 1992, OXFORD, GB, pages 3863-3866,
XP002038405

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The principal objects of the present invention are :


- 5 - novel organoselenium compounds : aromatic diselenides and selenosulphides;
- the use of said novel compounds as antioxidant ;
- pharmaceutical compositions containing them ;
- a method of preparation of said novel compounds.

10 STATE OF THE PRIOR ART:

[0002] In aerobic organisms, during the metabolism of oxygen, very reactive entities are generated whose accumulation causes deleterious effects. These organisms possess a system of regulation, composed of enzymes and small molecules which enable controlling the production of these reactive oxygen entities. Amongst the various components of this regulation system, often called Antioxidant Defence System, glutathione peroxidases play a central role in the prevention of «oxidative stress» and its deleterious consequences. These antioxidant and cytoprotecting enzymes enable degrading the endogenous or exogenous cytotoxic hydroperoxides.

[0003] These enzymes catalyse the reduction of hydrogen peroxide (reaction 1) or that of organic hydroperoxides (reaction 2) by reduced glutathione (GSH):

20 reaction 1 :



25 reaction 2 :



30 [0004] The active sites of these enzymes all contain one essential selenium atom in the form of a selenocysteine residue incorporated in the polypeptide chain.

[0005] The selenium is incorporated from selenite salts, or selenate salts, or L-seleno-methionine salts arising from food (see C.K. Chow and J. Jeng; in *Selenium in Medicine and Biology*, M.D. Spallholz and H.E. Ganther Eds.; (1986); Academic Press). In situations of selenium deficiency in the diet, the concentrations and activities of these glutathione peroxidases gradually decrease (see Y.X. Wang and J. Kiem; *Biological Trace Elements Res.*; (1988); 15; 89 and see R. Reiter and A.Wendel; *Biochem.Pharmacol.*; (1983); 32; 3063-3067); this leads to an acute susceptibility to the oxidative stress (see D.B.Coursin and H.P.Cihla, Thorax, (1996), 51. 479-483). The provision of selenium in the diet is therefore a limiting factor in the biosynthesis of glutathione peroxidases.

40 [0006] The protecting role of glutathione peroxidases, in situations wherein the production of hydroperoxides rises, has been demonstrated following experiments of direct intracellular micro-injection of the enzyme of erythrocyte origin which have enabled demonstrating its cytoprotective effect upon the viability of fibroblasts or endothelial cells exposed to an oxidative stress (see C. Michiels *et al.*; *Experiment. Cell Res.*; (1988); 179; 581-589). On the other hand, it has been shown that the survival of human fibroblasts is appreciably lowered when glutathione peroxidase is inhibited (see C. Michiels *et al.*, *J.Eur.Biochem.*; 1988, 177, 435-441).

45 [0007] Furthermore, the glutathione peroxidases are themselves particularly sensitive to an over-production of hydroperoxides and are rapidly inhibited under these conditions (see H. Ochi *et al.*, *Arch.of Biochem.Biophys.*, (1992), 294, 2, 407-411).

50 [0008] A certain number of pathologies such as certain ischaemic cardiomyopathies for example (see J.Chaudière, in *Biologie des lipides chez l'Homme*, L.Doustes-Blazy and F.Mendy eds, Edition Médicale Internationale - Paris, (1988), 137-154 and see D. Vitoux *et al.*; *Ann.Biol.Clin.*; (1996); 54; 5; 181-187) are associated with a lowering of glutathione peroxidase activity.

55 [0009] The demonstration of the essential role of the selenium at the active centre of these ubiquitous enzymes (see J.T.Rotruck, in *Selenium in Biology and Medicine*, Spallholz, J.E. Martin, J.L. Ganther H.I. eds., Avi Publishing Co, Westport, (1981), 10-16) as well as the importance of the selenium in the regulation of oxidative damages generated during certain pathologies (see Cadenas, E. and Sies, H., *Adv.Enz. Regul.*, (1985), 23, 217-237 and see Ursini, F., Bindoli, A., *Chem. Phys.Lipids*, (1987), 44, 255-276) has enabled the emergence of a novel class of organoselenium compounds as potential drugs.

[0010] Two types of compounds have been designed and prepared to this end.

[0011] On one side, modified macromolecules possessing a selenium atom introduced chemically such as seleno-subtilisin (see Z.P.Wu and D.Hilvert, *J.Am.Chem.Soc.*, (1990), 112, 5647-5648) or even a seleno-abzyme (see G.M. Luo *et al.*, *Biochem.Biophys.Res.Comm.*, 1994, 198, 3, 1240-1247). However, the use of proteins with a therapeutic aim is difficult to envisage for the following reasons:

- * their biostability is often insufficient;
- * an efficient method for ensuring their intra-cellular targeting does not exist;
- * they cannot be administered orally.

[0012] On the other side, synthetic molecules of low molecular weight have been synthesized, of which 2-phenyl-1,2-benzisoselenazolin-3-one (ebselen) (see H.Sies, *Free Rad.Biol.Med.*, (1993), 14, 313-323) was the first compound described as having a glutathione peroxidase activity. Homologs of said ebselen, i.e. 2H-3,4-dihydro-1,2-benzisoselenazin-3-ones, have also been described by Pierre V. Jacquemin *et al.* in *Tetrahedron Letters*, (1992), Vol. 33, No. 27, 3863-3866. In fact, several organoselenium derivatives have been described as glutathione peroxidase mimics, i. e. capable of reducing hydroperoxides in the presence of a biological thiol such as glutathione or lipoic acid (see I.A. Cotgreave *et al.*, *Biochem.Pharmacol.*, (1992), 43, 793-802 and C.M.Andersson *et al.*, *Free Rad.Biol.Med.*, (1994), 16, 17-28 and S.R.Wilson *et al.*, *J.Am.Chem.Soc.*, (1989), 111, 5936-5939 and V.Galet *et al.*, *J.Med.Chem.*, (1994), 37, 2903-2911). The patent application WO-A-95/27706 describes compounds of benzisoselenazoline and benzisose-lenazine structure having a glutathione peroxidase activity *inter alia*. US patents 5,128,365 and 5,321,138 themselves describe organic diselenides having glutathione peroxidase activity.

[0013] These organoselenium compounds, which are mimics of glutathione peroxidase, invariably produce catalytic intermediates of the selenol and/or diselenide type.

[0014] Amongst these, 2-phenyl-1,2-benzisoselenazolin-3-one (ebselen) and some of its derivatives do not seem to have any major toxic effect (see A. Wendel *et al.*; *Biochem. Pharmacol.*; (1984); 33; 3241-3245 and S.D. Mercurio and G.F. Combs; *Biochem. Pharmacol.*; (1986); 35; 4505-4509). 2-Phenyl-1,2-benzisoselenazolin-3-one (ebselen) is however very little soluble in water, even in the presence of an excess of glutathione GSH, which limits its pharmacological applications.

[0015] The biochemical and pharmacological properties of the organoselenium compounds which have been synthesised and studied have been recently reviewed (see M.J. Parnham and E. Graf; *Progress in Drug Res.*; (1991); 36; 9-47 and M.J.Parnham, *Exp.Opin.Invest.Drugs*, (1996), 5, 7, 861-570).

[0016] One of the aims of the present invention is to design organoselenium compounds having a catalytic activity of the glutathione peroxidase type in the presence of physiological concentrations of glutathione GSH.

[0017] These compounds must be able to penetrate the target tissues or cells, be soluble in water at active concentrations and must not efficiently reduce oxygen into toxic by-products.

[0018] These aims are attained by virtue of the present invention which resides on the design of cyclic organoselenium compounds whose antioxidant and cytoprotecting activities have been demonstrated by the Applicant and which are given below.

[0019] From a chemical point of view, very few 4(5)-seleno-imidazole derivatives have been described in the literature. Generally, these 4(5)-seleno-imidazole derivatives have been accessed according to 2 main routes, namely:

- * either by reaction between a derivative of type N-(trialkylsilyl)-imidazole and an arylselenium halide (see T.G. BACK and R.G. KERR; *Can. J.Chem.*; (1986); 64; 2; pages 308-301);
- * or by nucleophilic substitution between a 4-halo-imidazole derivative and a selenium derivative such as selenourea (see G.H. MILNE and L.R. TOWNSEND; *J.Carbohydr.Nucleosides, Nucléotides*; (1976); 3; 3; pages 177-183) or sodium hydrogen selenide (see G.H. MILNE and L.R. TOWNSEND; *J.Heterocycl.Chem.*; (1976); 13; 4; pages 745-748);

[0020] One of the objects of the invention described in this patent is to propose a novel method of introducing selenium in position 4(5) of an imidazole ring by reaction with an electrophilic selenium derivative such as selenium (Se⁰) chloride.

DESCRIPTION OF THE INVENTION:


[0021] The aim of the present invention is:

- 1) to solve the novel technical problem consisting of providing novel aromatic diselenides and selenosulphides having a very good antioxidant and cytoprotecting activity, thus constituting valuable active principles of pharmaceutical compositions;

2) to solve the novel technical problem above according to a solution which includes a method of preparation of these novel compounds which is easy to carry out.

[0022] The technical problems set forth above are solved for the first time in a simultaneous manner by the present invention in a simple way; the method of preparation of said novel compounds being relatively easy to carry out and giving good yields.

[0023] According to its first aspect, the present invention thus relates to novel organoselenium compounds - aromatic diselenides and selenosulphides - having the general formula (I) below :



in which :

R = hydrogen ; -C(R₁R₂)-A-B


R₁ = lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl;

R₂ = lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl;

A = CO ; (CR₃R₄)_n ;

B represents NR₅R₆ ; N⁺R₅R₆R₇Y⁻ ; OR₅ ; SR₅ ;


Ar = an optionally substituted phenyl group or an optionally substituted radical of formula :



in which Z represents O ; S ; NR₅ ; when R = -C(R₁R₂)-A-B


or

Ar = a radical of formula



in which Z = O ; S ; NR₅ ; when R is hydrogen ;


X = Ar(R)-Se- ; -S-glutathione ; -S-N-acetylcysteine ; -S-cysteine ; -S-penicillamine ; -S-albumin ; -S-glucose ;



25

30

R_3 = hydrogen ; lower alkyl ; optionally substituted aryl, optionally substituted lower aralkyl ;
 R_4 = hydrogen ; lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl ;
 R_5 = hydrogen ; lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted
heteroaryl ; optionally substituted lower heteroaralkyl ; CO(lower alkyl) ; CO(aryl) ; SO_2 (lower alkyl) ; SO_2 (aryl) ;
 R_6 = hydrogen ; lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted
heteroaryl ; optionally substituted lower heteroaralkyl ;
 R_7 = hydrogen ; lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted
heteroaryl ; optionally substituted lower heteroaralkyl ;
 R_8 = hydrogen ; lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted
heteroaryl ; optionally substituted lower heteroaralkyl ; trifluoromethyl ;



$n = 0$ or 1 ;

X^+ represents the cation of a pharmaceutically acceptable base ;

Y^- represents the anion of a pharmaceutically acceptable acid ;

55 and their salts of pharmaceutically acceptable acids or bases ;
with the provisos that :

when

$R = -C(R_1R_2)-(CR_3R_4)-B$ with $B = NR_5R_6$ or $N^+R_5R_6R_7Y^-$
 and
 $X = Ar(R)-Se-$ with Ar = optionally substituted phenyl,

5 then
 $-C(R_1R_2)$ is different from (CR_3R_4) ;
 and
 when

10 Ar = phenyl and
 $R = -C(R_1R_2)-C(O)-B$ with $B = NH_2$ or $NHCH_3$ or $NHCH_2C_6H_5$ or NHC_6H_5
 and
 $X = Ar(R)-Se-$,

15 then R_1 and R_2 cannot simultaneously represent a methyl group.

[0024] Said general formula (I) more particularly encompasses the 19 compounds, preparation of which is illustrated in the examples joined to this specification.

[0025] Said general formula (I) includes every stereoisomer, epimer and diastereoisomer, as a mixture or in isolated form.

20 [0026] It also includes, as indicated, the salts of pharmaceutically acceptable acids or bases of said compounds of formula (I).

[0027] Amongst the pharmaceutically acceptable acids, hydrochloric, hydrobromic, hydroiodic, sulphuric, tartaric, methanesulphonic, trifluoromethane-sulphonic acid, ... can be cited in a non-limiting way.

25 [0028] Amongst the pharmaceutically acceptable bases, sodium hydroxide, potassium hydroxide, alkali metal or alkaline earth metal carbonates, or organic bases such as triethylamine or arginine, ... can be cited in a non-limiting way.

[0029] Within the context of the present description and annexed claims :

- the terms "lower alkyl and lower alkoxy (see below)" are understood as meaning straight or branched alkyl and alkoxy groups having from 1 to 8 carbon atoms ;
 - 30 - the term "aryl" is understood as meaning an aromatic group selected from phenyl and naphthyl groups ;
 - the term "heteroaryl" is understood as meaning a mono- or bicyclic aromatic group, each cycle, or ring, comprising five or six atoms and said cycle, or ring, or both cycles, or rings, including in its carbon skeleton from one to three heteroatoms selected from nitrogen, oxygen and sulphur ;
 - the terms "lower aralkyl" and "lower heteroaralkyl" are understood as meaning, in view of the definitions above, phenyl(C_1-C_8)alkyl or naphthyl(C_1-C_8)alkyl and heteroar(C_1-C_8)alkyl respectively;
 - 35 - the term "substituted" concerning the terms aryl, aralkyl, phenyl, radical (five-membered, including Z), heteroaryl, heteroaralkyl, as defined above, signifies that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C_1-C_8)alkyl, trifluoromethyl, (C_1-C_8)alkoxy, hydroxy, nitro, amino, (C_1-C_8)alkylamino, di(C_1-C_8)alkylamino, sulphonyl, sulphonyl, sulphonamide, sulphonyl(C_1-C_8)alkyl,
 - 40 carboxyl, carbalkoxyl, carbamide (it being possible for said (C_1-C_8)alkyl groups to be linear or branched) or substituted with one or more halogen atoms;
 - the term aminoacyl, which concerns the glutathionyl, cysteinyl, N-acetylcysteinyl or even the penicillaminyl group in the definition of X, signifies any natural aminoacid such as alanine, and leucine, ... for example.
- 45 when R_3 and/or R_6 represents a hydrogen atom, the invention also covers the salts obtained with the pharmaceutically acceptable acids.

[0030] Said novel compounds have proved to be, as specified above, excellent antioxidant agents, the use of which is recommended by the Applicant in various fields. This use of said novel cyclic organoselenium compounds of the invention - compounds having formula (I) as defined above - as antioxidant agents, constitutes the second aspect of said invention.

50 [0031] Within the context of this second aspect, the use of said compounds of formula (I), is more particularly claimed as antioxidant :

- intended to be added to preserving media of grafts for transplantation of organs of human or animal origin such as the heart, the liver, the kidney and the lungs ;

and

- intended for (as active principle) the manufacture of pharmaceutical compositions with antioxidant activity, suitable especially :

- * for treatments of any physiopathological condition in which an over-production of cytotoxic hydroperoxides contributes to the functional impairments of cells or tissues ; and more particularly including :
- * the treatment of inflammatory and/or ischaemic cardio- and cerebro-vascular pathologies, such as the preventive and/or curative treatment of arterial restenoses following an angioplasty, the preventive and/or curative treatment of arterial stenoses following artery allografts, the treatment of intermittent claudication in patients affected with obstructive ischaemia of the lower members, the treatment of cerebro-vascular accidents of ischaemic origin ;
- * the treatment of inflammatory and/or ischaemic digestive pathologies, such as the treatment of acute inflammations of the bowel (Crohn's disease, hemorrhagic rectocolitis) ;
- * the treatment of inflammatory and/or ischaemic respiratory pathologies, such as the treatment of adult respiratory distress syndrome (ARDS) and infant respiratory distress syndrome (IRDS) ;
- * the treatment of inflammatory and/or ischaemic ophthalmic pathologies, such as the treatment of glaucoma ;
- * the treatment of cataracts ;
- * the treatment of acute ophthalmic allergies ;
- * the treatment of impairments of the retina which are associated with a macular degeneration ;
- * the treatment of viral infections causing an immuno-deficiency, such as the treatment of AIDS ;
- * the treatment of post-radiotherapy fibroses.

[0032] Generally, the potential therapeutic applications of the compounds of the invention include the treatment of any physiopathological condition in which an over-production of cytotoxic hydroperoxides contributes to the functional impairments of cells or tissues. Such an over-production of hydroperoxides can be endogenous and secondary to the activation of the intra-cellular metabolic pathways such as, for example, those of the flavine or cytochrome P-450 oxygenases, those of the lipoxygenases, those of the monoamine oxidases. The over-production can also be due to the activation of the endothelial cells (xanthine oxidase, 15-lipoxygenase), or of blood platelets (cyclooxygenase and 12-lipoxygenase). It can also be due to the activation, by cytokines such as TNF- α for example, of inflammatory and/or immune cells such as neutrophils, macrophages or lymphocytes for example. It may also be due to an intoxication by a free-radical generating xenobiotic. Finally, it may be due to a voluntary irradiation such as practised during a radiotherapy, or an accidental irradiation.

[0033] More particularly, the second aspect of the present invention includes the use of compounds of the invention for the manufacture of pharmaceutical compositions intended for the treatment :

- * of inflammatory diseases of the bowel such as Crohn's disease or hemorrhagic rectocolitis ;
- * of adult respiratory distress syndrome and infant respiratory distress syndrome ;
- * of cataracts ;
- * of AIDS ;
- * of post-radiotherapy fibroses.

[0034] From the second aspect of the present invention - use of the novel compounds de formula (I) as antioxidant agents -, such as described above, comes the third aspect which is dealt with now, namely the pharmaceutical compositions containing said compounds of formula (I) as active principle.

[0035] Thus, according to its third aspect, the present invention relates to pharmaceutical compositions, notably having an antioxidant activity, and comprising at least one organoselenium compound of the general formula (I), or one of its pharmaceutically acceptable salts of an acid or a base, as active ingredient, optionally incorporated in a pharmaceutically acceptable excipient, carrier or vehicle.

[0036] Said pharmaceutical compositions of the invention, according to an advantageous embodiment, contain said active ingredient in an amount between 0.1 and 5% by weight, advantageously between 0.1 and 1 % by weight based on their total weight. According to another advantageous embodiment, said compositions are in the form of unit doses comprising from 1 to 500 mg of at least one cyclic organoselenium compound of the invention (optionally incorporated in a pharmaceutically acceptable excipient, carrier or vehicle).

[0037] The pharmaceutical compositions of the invention can be formulated for, or intended for, oral, rectal or topical administration, (the compounds of formula (I) may especially be formulated for ophthalmic applications in the form of an eye lotion) or even as intra-ventricular, intra-muscular, subcutaneous or intravenous injections.

[0038] The pharmaceutically acceptable excipients, vehicles and carriers which can be included in their formulation are products which are well-known to the person skilled in the art and are not described in detail here.

[0039] The pharmaceutical compositions of the invention which contain the antioxidant agents disclosed by the

present invention (compounds of formula (I)) are especially suitable for the treatment of any physiopathological condition in which an over-production of cytotoxic hydroperoxides contributes to the functional impairments of cells or tissues ; it being possible for said over-production of hydroperoxides to be due to any one of the causes presented above in the present description, with reference to the second aspect of the invention (activation of the intra-cellular metabolic pathways, enzyme activation, macrophage or lymphocyte activation, intoxication by a free-radical generating xenobiotic, voluntary or accidental irradiation).

[0040] More specifically, said pharmaceutical compositions are suitable for the treatment of the pathologies listed above in the present description (with reference to the second aspect of the invention).

[0041] It is hereby specified that the antioxidant and therapeutic or pharmacological activities of the cyclic organoselenium compounds of the general formula (I) above have been demonstrated according to safe and reliable tests well-known to the person skilled in the art, which comprise :

a/ measuring the glutathione peroxidase activity;

b/ measuring the cytoprotective effect in human umbilical vein endothelial cells.

[0042] It is hereby incidentally mentioned that the preparation of pharmaceutical compositions incorporating an effective amount of at least one organoselenium compound of formula (I) according to the invention as well as the therapeutic treatments implying the use of such a compound make up an integral part of the present invention.

[0043] According to its last aspect, given below, the invention even relates to a method of preparation of said organoselenium compounds of formula (I). In fact, two synthetic routes are recommended:

- a route B for the compounds of formula (I) in which $R = H$;
- a route A for the other compounds of formula (I) in which $R \neq H$; $R = -C(R_1R_2)-A-B$.

[0044] First of all, said route A is presented. It comprises the following essential steps :

a/ preparing or using an orthohalo(hetero)arylacetonitrile derivative, optionally mono-or gem- disubstituted in the benzylic position; then, according to the series considered :

- for the preparation of said compounds of formula (I) in which $A = (CR_3R_4)_n$ and $n = 0$ (A does not exist):

b1/ hydrolyzing said nitrile derivative into an amide derivative,

c1-1/ transforming this amide derivative into an amine derivative by a transposition reaction according to conventional methods,

d1-1/ allowing said amine derivative to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylisoselenazoline derivative,

e1-1/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, according to conventional procedures, said (hetero)arylisoselenazoline derivative ;

- for the preparation of said compounds of formula (I) in which $A = CO$:

b1/ hydrolyzing said nitrile derivative into an amide derivative,

c1-2/ allowing said amide derivative to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylisoselenazone derivative,

d1-2/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, said (hetero)arylisoselenazone derivative; - for the preparation of said compounds of formula (I) in which $A = CH_2$:

b2/ reducing said nitrile derivative into an amine derivative with the aid of borane for example in an ethereal solvent such as tetrahydrofuran for example,

c2/ allowing said amine derivative to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylisoselenazine derivative,

d2/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, according to conventional procedures, said (hetero)arylisoselenazine derivative;

- for the preparation of said compounds of formula (I) in which $A = (CR_3R_4) (\neq CH_2)$:

b3/ carrying out a mono- or a bis-C-alkylation of said nitrile derivative according to conventional methods, with the aid of an organolithium derivative for example, in an ethereal solvent such as tetrahydrofuran;

c3/ allowing the amine derivative obtained to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)aryliselenazine derivative;

d3/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, according to conventional procedures, said corresponding (hetero)aryliselenazine derivative ;

b/ reducing, in a polar solvent such as methanol, the cyclic compound obtained, according to any one of the above synthetic routes, with the aid of a metal hydride, such as sodium borohydride, intervening in an amount corresponding to a half-reducing equivalent ; for the preparation of said compounds of formula (I) in which $X = \text{Ar(R)-Se-}$;

or
allowing said cyclic compound obtained according to any one of the above synthetic routes to react at ambient temperature, with the thiol compound corresponding to the values of $X \neq \text{Ar(R)-Se-}$, for the preparation of said compounds of formula (I) in which $X \neq \text{Ar(R)-Se-}$.

[0045] Said method is close to that described in the application WO-A-95/27706. It consists, as appears clearly above, of two main steps :

- upon completion of the first (a/), cyclic organoselenium compounds are obtained ;
- during the second (b/), said cyclic organoselenium compounds are either reduced or allowed to react with a thiol.

[0046] The implementation of each of its steps does not give rise to any particular difficulty to the person skilled in the art.


[0047] According to the advantageous variants of implementation of said method:

- the nucleophilic selenium derivative (which intervenes in steps d1-1/, c1-2/, c2/ and c3/) is a selenocyanate salt, such as potassium selenocyanate for example, which can be :

- * either generated *in situ* from selenium metal Se(0) and a cyanide salt, such as potassium cyanide for example,
- * or added to the reaction medium as such ;

- the copper salt Cu(I) (which intervenes in the same steps) is cuprous iodide ;
- the polar organic salt (which intervenes in the same steps) is dimethylformamide.


[0048] Route B is now presented. This is an original method of preparation, notably of 4(5)-seleno-imidazole derivatives of formula (I), more specifically compounds of the invention of formula (II) below :



in which Z and R₈ (and X) are defined as above with reference to the general formula (I).

[0049] Said original method comprises the following essential steps :

a/ preparing or using a derivative of formula (III) :



N-alkylated, N-arylated, N-acylated or N-sulphonylated when $Z = NR_5$; then, according to the series considered :

- for the preparation of said diselenide compounds of formula (II) in which $R_8 = \text{hydrogen}$:
 - b1/ allowing said derivative of formula (III) to react with an electrophilic selenium derivative, in a non-polar solvent;
 - c1/ if necessary, de-acylating or de-sulphonylating, according to usual procedures, the compound obtained ;
- for the preparation of said diselenide compounds of formula (II) in which $R_8 = \text{lower alkyl, optionally substituted lower aralkyl, optionally substituted lower heteroaralkyl}$:
 - b2/ alkylating said derivative of formula (III) in position 2 in treating it firstly with an organolithium base such as lithium diisopropylamide, then, with a halide, notably a lower alkyl, optionally substituted lower aralkyl or optionally substituted lower heteroaralkyl iodide ;
 - c2/ allowing said alkyl derivative to react with an electrophilic selenium derivative, in a non-polar solvent;
 - d2/ if necessary, de-acylating or de-sulphonylating, according to usual procedures, the compound obtained ;
- for the preparation of said diselenide compounds of formula (II) in which $R_8 = \text{optionally substituted aryl, optionally substituted heteroaryl}$:
 - b3/ treating said derivative of formula (III), in a non-polar organic solvent, in the presence of a strong base, with a trialkyltin halide or a zinc halide in order to obtain the corresponding stannyl or zinc derivative then treating said stannyl or zinc derivative with a haloaromatic derivative, in the presence of palladium, in a non-polar organic solvent ;
 - c3/ allowing the compound obtained to react with an electrophilic selenium derivative, in a non-polar solvent ;
 - d3/ if necessary, de-acylating or de-sulphonylating, according to usual procedures, the compound obtained ;
- b/ if necessary, allowing said diselenide compound obtained according to any one of the above synthetic routes, in a polar solvent, with an adequate mercaptan ; in order to obtain the corresponding selenosulphide compound.

[0050] For the preparation of the compounds - diselenides and selenosulphides - of the invention of formula (I) in which $R = H$, the diselenides are generally prepared as indicated above which are optionally transformed into selenosulphides (see B set forth above).

[0051] The alkyl group of the imidazole derivatives ($Z = NR_5$) N-alkylated is advantageously an alkyl group having from 1 to 6 carbon atoms, notably a methyl group.

[0052] The acyl group of the imidazole derivatives ($Z = NR_5$) N-acylated is advantageously a pivaloyl or a benzoyl group.

[0053] The sulphonyl group of the imidazole derivatives ($Z = NR_5$) N-sulphonylated is advantageously a tosyl group or a N,N-dimethylsulphonamide group.

[0054] According to an embodiment of this method (preparation variant for the diselenide compounds of formula (II) in which R_8 includes an aryl group : step b3/), this is characterised in that the trialkyltin halide is preferably tributyltin chloride or trimethyltin chloride. According to another embodiment of this method (same variant), this is characterised in that the zinc halide is preferably zinc chloride.

[0055] According to another particular embodiment of this method (same variant, same step b3/), this is characterised in that the above-mentioned strong base can be an alkyllithium, such as butyllithium for example, or a lithium amide, such as lithium diisopropylamide for example.

[0056] According to another embodiment of this method (same variant, same step b3/), this is characterised in that the non-polar organic solvent is preferably an ethereal solvent such as tetrahydrofuran for example.

[0057] According to yet another particular embodiment of this method (same variant, same step b3/), this is characterised in that the above-mentioned haloaromatic derivative is a chloro- or bromo- or iodo-aromatic derivative such as, for example, bromobenzene or 4-chloropyridine.

[0058] According to still another particular embodiment of this method (all variants), this is characterised in that the above-mentioned electrophilic selenium derivative can be selenium (Se^I) chloride.

[0059] According to still another particular variant of this method, this (all variants) is characterised in that the above-mentioned non-polar solvent can be dichloromethane for example.

[0060] When Z = NR₅, the protected nitrogen is deprotected, when necessary in a conventional way in a polar solvent. Said polar solvent can be tetrahydrofuran or acetonitrile, for example.

[0061] Finally, the polar solvent intervening in step b/ is advantageously acetonitrile.

[0062] Other aims, characteristics and advantages of the invention will appear clearly in the light of the following explanatory description made with reference to various non-limiting Examples given solely as illustration and which in no way limit the scope of the invention. In the Examples, all percentages are given by weight unless otherwise indicated.

[0063] In the annexed figures,

- Figure 1 shows the inhibition of the TNF- α -induced interleukin 8 release by endothelial cells (HUVEC);
- Figure 2 shows the inhibition of the TNF- α -induced P-selectin expression by endothelial cells (HUVEC);
- Figure 3 shows the inhibition of the TNF- α -induced E-selectin expression by endothelial cells (HUVEC).

EXPERIMENTAL SECTION:

[0064] All reactions were carried out under an inert nitrogen atmosphere unless otherwise indicated.

[0065] Mass spectra were recorded on a Nermag R10-10B instrument. Ionisation used is either electron impact (EI) at 70 electron-volts or chemical ionisation (CI) in ammonia or isobutene, or fast atom bombardment (FAB) on a glycerol matrix.

[0066] The ¹H and ¹³C NMR spectra were recorded on a Varian Gemini-200 instrument, the ⁷⁷Se NMR spectra on a Bruker AMX 500 instrument. The chemical shifts are given in ppm with respect to tetramethylsilane (¹H and ¹³C NMR spectra) or to dimethylselenide (⁷⁷Se NMR spectra). The multiplicities are expressed as follows: "s" for singlet, "bs" for broad singlet, "d" for doublet, "t" for triplet, "q" for quadruplet and "m" for multiplet; "E" for even and "O" for odd.


[0067] The melting points (m.p. °C) were recorded on a Gallenkamp instrument and are given uncorrected.

[0068] Purification by liquid column chromatography was carried out, according to the case, with Merck® Si60 F₂₅₄ or basic aluminium oxide Merck® Al₂O₃ 90 (Activity I).

// Examples of synthesis of compounds of general formula I (\neq II):

Example 1: Preparation of di[2-[2'-(1'-amino-2'-methyl)propyl]phenyl]-diselenide:

[0069]



[0070] The 4,4-dimethylbenziselenazine derivative **BXT 51072** (5.65 g; 25 mmoles) is suspended in anhydrous methanol (10 ml). A preparation of this derivative has been described in Example 8 of the application WO-A-95/27706.

[0071] Sodium borohydride (260 mg; 6.84 mmoles) is added slowly to the reaction medium, at a temperature of 0°C; stirring is then continued for one hour at ambient temperature and under air. Tert-butyl methyl ether (40 ml) and a 50% saturated sodium chloride solution (10 ml) are added to the reaction medium, the latter is then decanted. The orange organic phase is washed with 10 ml of a 50% saturated sodium chloride solution, dried over magnesium sulphate and filtered.

[0072] The solvent is evaporated under reduced pressure. The desired product is obtained as an orange oil. (Beware: the product dissolved in organic solvents, such as, for example dichloromethane, slowly oxidises in air to give the starting material, **BXT 51072**.)

Yield: 95%

Physical characteristics:

* NMR ^1H : (CDCl_3)

1.41 ppm (s; 6H); 1.60 ppm (bs; 2H); 3.08 ppm (s; 2H); 7.00-7.35 ppm (m; 3H); 7.95 ppm (m; 1H).

[0073] This product contains 6% of BXT 51072 according to the comparative integration of the small supplementary signals at 1.26 ppm and 3.21 ppm.

* NMR ^{13}C : (CDCl_3)

27.82 ppm; 42.65 ppm; 52.21 ppm; 127.98 ppm; 128.22 ppm; 128.97 ppm; 130.47 ppm; 136.45 ppm; 146.87 ppm.

* NMR ^{77}Se : (CH_3SeCH_3)

493 ppm

* MS: (Cl, isobutane)


457 (MH^+ ; 55%); 228 (100%).

* HRMS:

calc. for $\text{C}_{20}\text{H}_{29}\text{N}_2\text{Se}_2$: 457.0661; exp.: 457.0665.

Example 2: Preparation of dif[2-[2'-(1'-amino-2'-methyl)propyl]phenyl]-diselenide dihydrochloride: BXT 51125

[0074]



[0075] The 4,4-dimethylbenzisoselenazine derivative BXT 51072 (27.12 g; 0.12 mol) (see Example 8 of WO-A-95/27706) is suspended in anhydrous methanol (50 ml). Sodium borohydride (1.25 g; 33 mmoles) is added in portions of 50 mg for 30 min. to the reaction medium at a temperature of 0°C ; stirring is then continued vigorously for 5 hours at ambient temperature and under air. Tert-butyl methyl ether (200 ml) and a 50% saturated sodium chloride solution (50 ml) are added to the reaction medium, the latter is then decanted. The orange organic phase is washed with 50 ml of a 50% saturated sodium chloride solution, dried over magnesium sulphate and filtered. This solution is added slowly (50 min.) to an ethanolic solution of hydrogen chloride (68 ml; 4.1M; 0.28 mol) at a temperature of 5°C . Stirring is continued vigorously for 30 min. The suspension obtained is filtered. The yellow precipitate is washed with 5x50 ml of tert-butyl methyl ether, then dried. The desired compound is obtained as a yellow powder.

Yield: 95%

Physical characteristics:

* m.p. $^\circ\text{C}$: 283°C (dec.)

* NMR ^1H : (D_2O)

1.39 ppm (s; 6H); 3.42 ppm (s; 2H); 7.06 ppm (m; 1H); 7.25 ppm (m; 1H); 7.36 ppm (m; 1H); 7.65 ppm (m; 1H).

* NMR ^1H : (CD_3OD)

1.56 ppm (s; 6H); 3.52 ppm (s; 2H); 7.21 ppm (td; 1H, $J = 8 - 8 - 1.5$ Hz); 7.36 ppm (td; 1H, $J = 8 - 8 - 1.5$ Hz); 7.41 ppm (dd; 1H, $J = 8 - 1.5$ Hz); 7.80 ppm (dd; 1H, $J = 8 - 1.5$ Hz).

* NMR ^{13}C : (D_2O)

24.95 ppm; 37.35 ppm; 46.29 ppm; 127.13 ppm; 127.27 ppm; 127.74 ppm; 127.90 ppm; 136.33 ppm; 142.07 ppm.


* microanalysis:

calc. for $\text{C}_{20}\text{H}_{30}\text{Cl}_2\text{N}_2\text{Se}_2$: C 45.56%; H 5.73%; N 5.31 %;

exp. C 45.50%; H 5.77%; N 5.44%.

Example 3: Preparation of di[2-[2'-(1'-ammonium-2'-methyl)propyl]phenyl]-diselenide di-paratoluenesulphonate: BXT 51108

[0076]



[0077] The 4,4-dimethylbenzisoselenazine derivative BXT 51072 (3.39 g; 15 mmoles) (see Example 8 of WO-A-95/27706) is dissolved in anhydrous methanol (75 ml). Sodium borohydride (627 mg; 16.5 mmoles) is added; then the reaction mixture is stirred for 30 min; at ambient temperature and under air. The methanol is evaporated under reduced pressure. The residus is taken up into 150 ml of ethyl acetate and washed with 2x150 ml of water; dried over magnesium sulphate and filtered. This solution is added to a solution of para-toluenesulphonic acid (3.13 g; 16.5 mmoles) in ethyl acetate (75 ml). The reaction mixture is brought to a temperature of 0°C, and stirring is continued for 30 min. The suspension obtained is filtered; The precipitate is washed with 4x25 ml of tert-butyl methyl ether, then dried. The desired compound is obtained as a yellow powder.


Yield: 89%

Physical characteristics:

- * m.p.°C: 176°C (dec.)
- * NMR ¹H: (CD₃OD)
1.54 ppm (s; 6H); 2.35 ppm (s; 3H); 3.49 ppm (s; 2H); 7.10-7.47 ppm (m; 3H); 7.21 ppm (d; 2H, J = 8 Hz); 7.69 ppm (d; 2H, J = 8 Hz); 7.78 ppm (m; 1H).
- * NMR ¹³C: (CD₃OD)
21.31 ppm; 27.58 ppm; 40.75 ppm; 49.64 ppm; 127.37 ppm; 129.72 ppm; 130.00 ppm; 130.14 ppm; 130.56 ppm; 131.03 ppm; 139.22 ppm; 142.08 ppm; 143.76 ppm; 145.55 ppm.
- * MS: (FAB⁺, para-nitrobenzyl alcohol)
457/455 (M⁺⁺-H⁺, 67%); 230 (76%); 228 (100%); 226 (53%); 212 (38%); 197 (31%); 154 (48%); 136 (45%).
- * HRMS:
calc. for C₂₀H₂₉N₂Se₂ (M⁺⁺-H⁺): 457.0661. exp.: 457.0665.

Example 4: Preparation of di[2-[2'-(1'-amino-2'-methyl)propyl]-4-methoxy]phenyl-diselenide:

[0078]



[0079] The 4,4-dimethyl-6-methoxybenzisoselenazine derivative BXT 51077 (256 mg; 1 mmol) is dissolved in anhydrous methanol (5 ml). A preparation of this compound has been described in Example 11 of the application WO-A-95/27706. Sodium borohydride (9.5 mg; 0.25 mmoles) is added; then the reaction mixture is stirred for 30 min; at ambient temperature and under air. The methanol is evaporated under reduced pressure. The residue is taken up into 20 ml of water, then extracted with 2x20 ml of dichloromethane. The organic phases are combined, washed with 20 ml of a saturated sodium chloride solution, dried over sodium sulphate and filtered. The solvent is evaporated off under reduced pressure. The desired compound is obtained as an orange oil, in a mixture with 3% of starting material, BXT 51077.

Yield: 82%

(Beware: the product dissolved in organic solvents, such as, for example dichloromethane, slowly oxidises in air to give the starting material, BXT 51077.)

Physical characteristics:


- * NMR ^1H : (CDCl_3)
0.8 ppm (bs; 2H); 1.34 ppm (s; 6H); 3.06 ppm (s; 2H); 3.74 ppm (s; 3H); 6.60 ppm (dd; 1H, $J = 8.5 - 2.5$ Hz); 6.84 ppm (d; 1H, $J = 2.5$ Hz); 7.65 ppm (d; 1H, $J = 8.5$ Hz).

[0080] This product contains 3% of BXT 51077 according to the comparative integration of the small supplementary signal at 1.24 ppm.

- * NMR ^{13}C : (CDCl_3)
28.57 ppm; 42.77 ppm; 52.42 ppm; 55.42 ppm; 111.64 ppm; 116.26 ppm; 120.87 ppm; 140.25 ppm; 149.56 ppm; 160.11 ppm.
* NMR ^{77}Se : (CH_3SeCH_3)
521 ppm (signals of weak intensity at 694 ppm (corresponding to BXT 51077) and 345 ppm).
* MS: (CI; isobutane)
517 (MH^+ ; 27%); 258 (100%).

Example 5: Preparation of di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenide: BXT 51109

[0081]:



5A/ Preparation of 2,4,4-trimethyl-benzisoselenazine: BXT 51112

[0082] The 4,4-dimethylbenzisoselenazine derivative BXT 51072 (452 mg; 2 mmoles) (see Example 8 of WO-A-95/27706) is dissolved in 4 ml of tert-butyl methyl ether. Methyl para-toluenesulphonate (372 mg; 2 mmoles) is added, then the solution is refluxed for 14h. The reaction mixture, brought to ambient temperature, is diluted with tert-butyl methyl ether (20 ml), then extracted with hydrochloric acid (1N; 3x15 ml). The aqueous phases are combined. After making it alkaline (pH = 12), the aqueous phase is extracted with tert-butyl methyl ether (3x20 ml). The organic phases are combined, washed with 20 ml of a saturated sodium chloride solution, dried over sodium sulphate and filtered. The solvent is evaporated off under reduced pressure. The desired compound is obtained as a yellow oil after purification by liquid chromatography on a silica column (eluent: cyclohexane-ethyl acetate 9/1).

Yield: 47%

Physical characteristics:

- * NMR ^1H : (CDCl_3)
1.23 ppm (bs; 6H); 2.78 ppm (s; 3H); 3.06 ppm (s; 2H); 7.0-7.13 ppm (m; 3H); 7.37-7.46 ppm (m; 1H).
* NMR ^{13}C : (CDCl_3)
29.85 ppm; 37.05 ppm; 48.51 ppm; 72.11 ppm; 126.09 ppm; 126.30 ppm; 126.85 ppm; 127.85 ppm; 129.69 ppm; 142.64 ppm.
* MS: (EI, 70 eV)
241 (M^+ ; 100%); 198 (62%), 183 (76%), 115 (40%).

5B/ Preparation of di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenide: BXT 51109

[0083] The preceding derivative BXT 51112 (140 mg; 0.58 mmol) is dissolved in anhydrous methanol (5 ml). Sodium borohydride (5.5 mg; 0.145 mmole) is added; then the reaction mixture is stirred for 30 min., at ambient temperature and under air. The methanol is evaporated under reduced pressure. The residue is taken up into 20 ml of hydrochloric acid (1N), then washed with 3x10 ml of tert-butyl methyl ether. The aqueous phase is made alkaline, then extracted with 3x20 ml of dichloromethane. The organic phases are combined, washed with 20 ml of a saturated sodium chloride solution, dried over sodium sulphate and filtered. The solvent is evaporated under reduced pressure. The desired

compound is obtained as an orange oil.

(Beware, the product dissolved in organic solvents, such as, for example dichloromethane, slowly oxidises to give the starting material BXT 51112.)


Yield: 75%

Physical characteristics:

- * NMR ^1H : (CDCl_3)
1.41 ppm (s; 6H); 2.05 ppm (bs; 1H); 2.23 ppm (s; 3H); 3.02 ppm (s; 2H); 7.0-7.20 ppm (m; 2H); 7.83 ppm (dd; 1H, $J = 8 - 1.5$ Hz); 7.83 ppm (dd; 1H, $J = 8 - 1.5$ Hz).
- * NMR ^{13}C : (CDCl_3)
28.70 ppm; 37.27 ppm; 41.28 ppm; 62.27 ppm; 127.91 ppm; 128.18 ppm; 129.54 ppm; 130.36 ppm; 136.37 ppm; 146.68 ppm.
- * MS: (CI, isobutane)
485 (MH^+ ; 68%); 242 (100%).

Example 6: Preparation of di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenide dihydrochloride: BXT 51130

[0084]



[0085] This compound is obtained, according to a method very similar to that of Example 2 from the preceding derivative (BXT 51112) as a yellow-orange powder.


Yield: 89%

Physical characteristics:

- * m.p.: 260°C (dec.)
- * NMR ^1H : (CD_3OD)
1.57 ppm (s; 6H); 2.56 ppm (s; 3H); 3.57 ppm (s; 2H); 7.24 ppm (td; 1H, $J = 8 - 8 - 1.5$ Hz); 7.37 ppm (td; 1H, $J = 8 - 8 - 1.5$ Hz); 7.47 ppm (dd; 1H, $J = 8 - 1.5$ Hz); 7.85 ppm (dd; 1H, $J = 8 - 1.5$ Hz).
- * NMR ^{13}C : (CD_3OD)
27.98 ppm; 35.20 ppm; 40.76 ppm; 59.66 ppm; 129.56 ppm; 130.18 ppm; 130.76 ppm; 131.29 ppm; 139.90 ppm; 145.56 ppm.

Example 7: Preparation of di[2-[2'-(1'-dimethylamino-2'-methyl)propyl]-phenyl]-diselenide: BXT 51110

[0086]



[0087] The derivative BXT 51109 obtained in situ, by the method of Example 5/B from BXT 51112 (360 mg; 1.5 mmoles) is dissolved under nitrogen in tert-butyl methyl ether (6 ml). Methyl para-toluenesulphonate (335 mg; 1.8 mmoles) is added, stirring is then continued for 24h at ambient temperature. Sodium hydroxide solution (1N; 20 ml) is added, then the mixture is extracted with 2x20 ml of dichloromethane. The organic phases are combined, washed with 20 ml of a saturated sodium chloride solution, dried over sodium sulphate and filtered. The solvent is evaporated under reduced pressure. The desired product is obtained as an orange oil after purification by liquid chromatography on a silica column (eluent gradient: cyclohexane-ethyl acetate 20/1 then methanol/triethylamine 100/1).


Yield: 69 %

Physical characteristics:

- * NMR ^1H : (CD_3OD)
1.42 ppm (s; 6H); 2.05 ppm (s; 6H); 2.7 ppm (bs; 2H); 7.02 ppm (td; 1H, $J = 7.5 - 7.5 - 1.5$ Hz); 7.14 ppm (td; 1H, $J = 7.5 - 7.5 - 1.5$ Hz); 7.33 ppm (dd; 1H, $J = 7.5 - 1.5$ Hz); 7.89 ppm (dd; 1H, $J = 7.5 - 1.5$ Hz).
- * NMR ^{13}C : (CDCl_3)
28.62 ppm; 41.53 ppm; 48.01 ppm; 70.51 ppm; 127.32 ppm; 127.50 ppm; 133.92 ppm; 137.14 ppm; 146.85 ppm.
- * MS: (FAB $^+$; para-nitrobenzyl alcohol)
513 (MH^+ ; 20); 256 (100%); 154 (50%); 136 (40%).

Example 8: Preparation of di[2-[2'-(1'-trimethylammonium-2'-methyl)propyl]-phenyl]-diselenide di-paratoluenesulphonate: BXT 51111

[0088]



[0089] The preceding derivative (102 mg; 0.2 mmol) is dissolved in tert-butyl methyl ether (4 ml). Methyl para-toluenesulphonate (93 mg; 0.5 mmol) is added, then the mixture is refluxed. After 24h, the reaction mixture, brought to ambient temperature, is diluted with tert-butyl methyl ether (20 ml), then extracted with water (3x15 ml). The aqueous phases are combined and washed with tert-butyl methyl ether (10 ml). The water is evaporated under reduced pressure. The desired product is obtained first as an orange oil. For the crystallisation, the oil is taken up into ethanol (1 ml), then ethyl acetate is added (20 ml). The suspension obtained is filtered and the precipitate washed with 2x5 ml of tert-butyl methyl ether. Thus the desired product is obtained as a yellow powder.


Yield: 48%

Physical characteristics:

- * m.p. $^{\circ}\text{C}$: 203 $^{\circ}\text{C}$ (dec.)
- * NMR ^1H : (CD_3OD)
1.69 ppm (s; 6H); 2.35 ppm (s; 3H); 2.91 ppm (s; 9H); 4.10 ppm (s; 2H); 7.22 ppm (d; 2H, $J = 8.5$ Hz); 7.27 ppm (td; 1H, $J = 8 - 8 - 1.5$ Hz); 7.41 ppm (td; 1H, $J = 8 - 8 - 1.5$ Hz); 7.61 ppm (dd; 1H, $J = 8 - 1.5$ Hz); 7.69 ppm (d; 2H, $J = 8.5$ Hz); 7.89 ppm (dd; 1H, $J = 8 - 1.5$ Hz).
- * NMR ^{13}C : (CD_3OD)
21.35 ppm; 31.01 ppm; 42.89 ppm; 56.27 ppm; 74.92 ppm; 127.31 ppm; 130.20 ppm; 130.35 ppm; 130.56 ppm; 131.25 ppm; 131.45 ppm; 139.70 ppm; 142.09 ppm; 144.04 ppm; 145.69 ppm.
- * MS: (FAB $^+$, glycerol)
713 ($(\text{M} - \text{Tos})^+$; 30%); 271 ($\text{M}^{++} - 2\text{Tos}$; 50%); 256 (80%); 212 (100%); 197 (85%); 130 (100%); 91 (50%).

Example 9: Preparation of di [3-[2'-(1'-amino -2'-methyl)propyl]-2-thienyl] diselenide : BXT 51099

[0090]



A/ Preparation of 2-bromo-3-bromomethyl-thiophene :

[0091] This compound is prepared from 2-bromo-3-methyl-thiophene (1.85 g; 10.4 mmoles) according to a method identical to that described in the literature by E. Campaigne and W.M. Lesuer (J.Am.Chem.Soc., 1949, 71, 333-335), and obtained as a colourless oil (2.45 g). The product is used as such in the next step. Yield : 95 %

Physical characteristics :

- * NMR ^1H : (CDCl_3)
4.43 ppm (s, 2H); 6.98 ppm (d, 1H, $J=5.7$ Hz); 7.24 ppm (d, 1H, $J=5.7$ Hz).

B/ Preparation of 3-(2-bromo)-thienyl- acetonitrile :

[0092] A round-bottomed flask, containing the preceding bromine derivative (2.10 g ; 8.2 mmoles) dissolved in methanol (8 ml), is cooled in a water bath. Under an inert atmosphere, sodium cyanide (0.610 g; 12.3 mmoles) is added all at once. The mixture is left for 4 hours, at ambient temperature. The solvent is evaporated under reduced pressure. The residue is taken up into 15 ml of water and extracted with 2 x 15 ml of dichloromethane. The organic phases are combined, washed with water, then with a saturated sodium chloride solution, dried over magnesium sulphate, then filtered. After evaporation of the solvent and chromatography on a silica column (eluent gradient pure cyclohexane, then cyclohexane/ethyl acetate (90/10)), the desired product is obtained as a pale yellow oil (1.05 g).

Yield : 64 %

Physical characteristics :

- * NMR ^1H : (CDCl_3)
3.64 ppm (s, 2H); 7.01 ppm (d, 1H, $J=5.8$ Hz); 7.31 ppm (d, 1H, $J=5.8$ Hz).
- * NMR ^{13}C : (CDCl_3)
18.62 ppm (E); 112.28 ppm (E); 117.12 ppm (E); 127.50 ppm (O); 128.03 ppm (O); 129.80 ppm (P).
- * MS: (EI; 70eV)
203/201 (M^{++} ; 45); 122 (M-Br; 100); 98(60).

C/ Preparation of 2-[3'-(2'-bromo)-thienyl]-2-methyl-propionitrile :

[0093] To a suspension of NaH (0.140 g; 4 mmoles) in anhydrous DMF (1 ml), kept at -10°C under an inert atmosphere, is added slowly (~ 5 -10 minutes), a solution containing the preceding derivative (0.202 g ; 1 mmole) and methyl iodide (0.570 g, 4 mmoles) in anhydrous DMF (1.5 ml). The reaction mixture is added at 0°C for one hour, then at ambient temperature for about 2 hours. The residue is taken up into 10 ml of water and extracted with 2 x 10 ml of ethyl acetate. The organic phases are combined, washed with 3 x 10 ml of water, then with a saturated sodium chloride solution, dried over magnesium sulphate, then filtered. The desired product is obtained, after evaporation of the solvent, as a yellow brown oil (0.240 g). It will be used as such in the next step.

Yield : \sim quantitative

Physical characteristics :

- * NMR ^1H : (CDCl_3)
1.80 ppm (s, 6H); 6.94 ppm (d, 1H, $J=5.9$ Hz); 7.24 ppm (d, 1H, $J=5.9$ Hz).
- * NMR ^{13}C : (CDCl_3)
27.35 ppm; 33.68 ppm; 109.74 ppm; 122.97 ppm; 126.39 ppm; 126.54 ppm; 138.32 ppm.
- * MS: (EI; 70eV)
231/229 (M^{++} ; 30); 216/214 (M-15; 70); 153/151 (M-Br; 45); 125/123 (42); 97 (100).

D/ Preparation of 2-[5'-(2'-bromo)-thienyl]-2-methyl-propionitrile :

[0094] The derivative, prepared in Example 9/C, (0.230 g ; 1 mmole) is dissolved in anhydrous THF (1 ml), under nitrogen. A solution of aluminium hydride AlH_3 in anhydrous THF (1.5 M; 1.0 ml; 1.5 mmoles) is added slowly to the reaction mixture at ambient temperature. The mixture is refluxed for 2 hours. Brought to ambient temperature, the medium is first hydrolysed, with 1 to 2 ml of water, then with 2 ml of a 2N solution of HCl. After evaporation of solvents, the residue is taken up into 5 ml of a 2N solution of HCl and extracted with 3 x 10 ml of TBME. The aqueous phase is made alkaline ($\text{pH}=12$), then extracted with 3 x 6 ml of ethyl acetate. The organic phases are combined, washed with 2 x 10 ml of water, then with a saturated sodium chloride solution, dried over magnesium sulphate, then filtered. The desired product is obtained after evaporation of the solvent, as a yellow oil (0.125 g).

Yield : 53 %

Physical characteristics:

- * NMR ^1H : (CDCl_3)
1.12 ppm (bs, $-\text{NH}_2$); 1.38 ppm (s, 6H); 2.98 ppm (s, 2H); 6.84 ppm (d, 1H, $J=5.8$ Hz); 7.16 ppm (d, 1H, $J=5.8$ Hz).

- * NMR ^{13}C : (CDCl_3)
26.38 ppm; 40.15 ppm; 52.19 ppm; 107.04 ppm; 125.61 ppm; 129.36 ppm; 145.17 ppm.
- * MS: (CI; isobutane)
236/234 (MH^+ ; 100); 220/218 ($\text{M}-\text{NH}_2$; 5); 154 ($\text{M}-\text{Br}$; 47); 125/123(12).

E/ Preparation of 4,4-dimethyl-thieno-[3,2-e]-isoselenazine :


[0095] To a solution of potassium selenocyanate (1.0 g ; 6.9 mmoles) in anhydrous DMF (7 ml), is added the derivative prepared as in Example 9/D (0.702 g; 3 mmoles). To this solution, cooled to 5°C and under an inert atmosphere, are then added copper I iodide (0.570 g ; 3 mmoles), then triethylamine (0.900 g ; 9 mmoles). The reaction mixture rapidly becomes deeply coloured, and is thus kept stirred for 17 hours, at ambient temperature and under nitrogen. 70 ml of an aqueous sodium cyanide solution are added (0.560 g; 11.4 mmoles) such that the copper iodide is complexed out, thus facilitating the two phases. After decantation, the aqueous phase is extracted with 70 ml of ethyl acetate. The organic phase is washed with 5 x 50 ml of water, then with 2 x 50 ml of a saturated sodium chloride solution, dried over magnesium sulphate, then filtered. The desired product is obtained after evaporation of the solvent and chromatography on a silica column (eluent cyclohexane/ethyl acetate (95/5)) as light yellow crystals (0.390 g).

Yield : 56%

Physical characteristics:

- * m.p. °C: 51.6-51.8 °C (hexane/ethyl acetate : 75/1)
- * NMR ^1H : (CDCl_3)
1.21 ppm (s, 6H); 3.18 ppm (s, 2H); 3.29 ppm (bs, -NH); 7.05 ppm (d, 1H, J=5.2 Hz); 7.27 ppm (d, 1H, J=5.2 Hz).
- * NMR ^{13}C : (CDCl_3)
28.63 ppm (O); 33.69 ppm (E); 61.50 ppm (E); 119.78 ppm (E); 123.93 ppm (O); 127.18 ppm (O); 141.84 ppm (P).
- * NMR ^{77}Se : (CH_3SeCH_3)
737.0 ppm.
- * MS: (EI; 70 eV)
233 (M^{+} ; 82); 204(79); 189(100); 124(22); 97(17).

[0096] The product obtained BXT 51097 is of the formula below :



F/ Said 4,4-dimethylthieno-[3,2-e]isoselenazine derivative BXT 51097 (0.114 g ; 0.50 mmoles), is dissolved in methanol (5 ml), at ambient temperature. To this solution cooled at 0°C, sodium borohydride is added (0.100 g; 2.5 mmoles). The reaction mixture is allowed to attain ambient temperature. After 15 to 20 minutes, the solvent is evaporated under reduced pressure. The residue, taken up into ethyl acetate, is washed 3-4 times with a saturated sodium chloride solution, then dried over magnesium sulphate and filtered. After evaporation of the solvent under reduced pressure, the desired product is obtained (BXT 51099), as a yellow oil (0.080 g).


Yield : 35%.

Physical characteristics:

- * NMR ^1H : (CDCl_3)
1.34 ppm (s, 6H); 2.92 ppm (s, 2H); 6.96 ppm (d, 1H, J=5.4 Hz); 7.36 ppm (d, 1H, J=5.4 Hz).
- * NMR ^{13}C : (CDCl_3)
27.78 ppm; 41.08 ppm; 53.96 ppm; 119.73 ppm; 129.95 ppm; 131.07 ppm; 152.48 ppm.
- * NMR ^{77}Se : (CH_3SeCH_3)
509.3 ppm
- * MS: (EI; 70 eV)
233 ($\text{M}^{+}/2$; 65); 204 (65); 189 (100); 124 (30); 97 (27).

Example 10: Preparation of di [2-[2'-(1'-amino-2'-methyl) propyl]-3-thienyl]-diselenide: BXT 51104

[0097]



A/ Preparation of 3-bromo-2-hydroxymethyl-thiophene:

[0098] 2-(3-bromothiophenyl)carboxylic acid (7 g; 34 mmoles) is dissolved in 25 ml of anhydrous THF. A solution of aluminium hydride AlH_3 in anhydrous THF (2 M; 42 ml; 84 mmoles) is added slowly at a temperature of 0°C . At the end of the addition, the reaction mixture is refluxed for 3h. After cooling to 0°C , water (200 ml), then hydrochloric acid (1N, 150 ml) are added. The mixture is decanted, and the aqueous phase extracted with 3 x 150 ml of tert-butyl methyl ether. The organic phases are combined, then washed with 150 ml of a saturated sodium chloride solution, dried over magnesium sulphate and filtered. The solvent is evaporated under reduced pressure. The desired product is obtained as a brown oil and is used as such for the next step.

Yield: 95%

Physical characteristics:

- * NMR ^1H : (CD_3OD)
4.69 ppm (s; 2H); 6.95 ppm (d; 1H, $J = 5.2$ Hz); 7.39 ppm (d; 1H, $J = 5.2$ Hz).
- * NMR ^{13}C : ($\text{DMSO}-d_6$)
57.80 ppm; 106.28 ppm; 126.11 ppm; 129.80 ppm; 141.15 ppm.
- * MS: (EI, 70 eV)
194/192 (M^{++} ; 80%); 177/175 (30%); 113 (50%); 98 (60%); 85 (100%).

B/ Preparation of 3-bromo-2-chloromethyl-thiophene:

[0099] The preceding derivative (6.2 g; 32 mmoles) is dissolved in anhydrous dichloromethane (180 ml). Thionyl chloride (5.7 g; 3.5 ml; 48 mmoles) is added slowly, then stirring is kept up for 18h at ambient temperature. The reaction mixture is poured into 200 ml of water, then decanted. The aqueous phase is extracted with 2 x 100 ml of tert-butyl methyl ether. The organic phases are combined, washed with a saturated sodium bicarbonate solution (200 ml), dried over magnesium sulphate and filtered. The solvent is evaporated off under reduced pressure. The desired product is obtained as a brown oil and is used as such for the next step.

Yield: 90 %

Physical characteristics:

- * NMR ^1H : (CDCl_3)
4.75 ppm (s; 2H); 6.95 ppm (d; 1H, $J = 5.2$ Hz); 7.20 ppm (d; 1H, $J = 5.2$ Hz).
- * NMR ^{13}C : (CDCl_3)
39.5 ppm; 112.5 ppm; 127.5 ppm; 131.0 ppm; 135.0 ppm.
- * MS: (EI, 70 eV)
212 (M^{++} ; 35%); 177 (100%); 96 (20%).

C/ Preparation of 2-(3-bromo)-thienyl-acetonitrile:

[0100] The preceding derivative (6 g; 28 mmoles) is dissolved in DMSO (100 ml). Sodium cyanide (2 g; 41 mmoles) is added, then stirring is kept up for 2h. The reaction mixture is poured into 200 ml of water. The mixture is extracted with 3 x 100 ml of tert-butyl methyl ether, then 2 x 100 ml of dichloromethane. The organic phases are combined, washed with 3 x 200 ml of water, dried over magnesium sulphate and filtered. The solvent is evaporated off under reduced pressure. The residue is taken up into 400 ml of cyclohexane and washed with 4 x 100 ml of water. The organic phase is dried over magnesium sulphate and filtered. The solvent is evaporated under reduced pressure. The desired product is obtained as a colourless oil after distillation of the residue in a Kugelrohr distillation apparatus ($T = 200^\circ\text{C}$,

p = 0.1 mbar).

Yield: 60%

Physical characteristics:

- 5 * NMR ^1H : (CDCl_3)
3.84 ppm (s; 2H); 7.00 ppm (d; 1H, J = 5.2 Hz); 7.31 ppm (d; 1H, J = 5.2 Hz).
* NMR ^{13}C : (CDCl_3)
18.39 ppm; 112.29 ppm; 116.30 ppm; 126.37 ppm; 126.65 ppm; 130.92 ppm.
* MS: (EI, 70 eV)
10 203/201 (M^{++} ; 29%); 122 (100%); 98 (40%).

D/ Preparation of 2-[2'-(3'-bromo-thienyl)]-2-methyl-propionitrile:

- 15 [0101] This compound is obtained according to a method very similar to that of Example 9/C from the preceding derivative as a yellow oil.

Yield: 90%

Physical characteristics:

- 20 * NMR ^1H : (acetone- d_6)
1.77 ppm (s; 6H); 7.02 ppm (d; 1H, J = 5.2 Hz); 7.44 ppm (d; 1H, J = 5.2 Hz).
* NMR ^{13}C : (CDCl_3)
20.11 ppm; 33.90 ppm; 109.68 ppm; 122.15 ppm; 124.39 ppm; 133.21 ppm; 137.27 ppm.
* MS: (EI, 70 eV)
25 231/229 (M^{++} ; 50%); 216/214 (100%); 189/187 (45%).

E/ Preparation of 2-[2'-(3'-bromo-thienyl)]-2-methyl-propylamine:

- 30 [0102] This compound is obtained according to a method very similar to that of Example 9/D from the preceding derivative as a pale yellow oil.

Yield: 80%

Physical characteristics:

- 35 * NMR ^1H : (CDCl_3)
1.10 ppm (bs; 2H); 1.44 ppm (s; 6H); 3.06 ppm (s; 2H); 6.93 ppm (d; 1H, J = 5.2 Hz); 7.07 ppm (d; 1H, J = 5.2 Hz).
* NMR ^{13}C : (CDCl_3)
26.45 ppm; 40.76 ppm; 51.78 ppm; 105.89 ppm; 123.01 ppm; 133.26 ppm; 145.00 ppm.
* MS: (CI, isobutane)
236/234 (MH^+ ; 40%); 154 (25%); 93 (100%).

- 40 F/ Preparation of 4,4-dimethyl-thieno-[2,3-e]-isoselenazine: BXT 51103


[0103] This compound is obtained according to a method very similar to that of Example 9/E from the preceding derivative as a yellow oil, which crystallises at -20°C .

Yield: 54%

Physical characteristics:

- 45 * m.p. $^\circ\text{C}$: 31°C
* NMR ^1H : (CDCl_3)
1.33 ppm (s; 6H); 3.20 ppm (s; 2H); 3.30 ppm (bs; 1H); 6.68 ppm (d; 1H, J = 5.3 Hz); 7.24 ppm (d; 1H, J = 5.3 Hz).
50 * NMR ^{13}C : (CDCl_3)
30.02 ppm; 34.16 ppm; 61.35 ppm; 119.64 ppm; 122.89 ppm; 124.57 ppm; 139.89 ppm.
* NMR ^{77}Se : (CH_3SeCH_3)
727.0 ppm.
* MS: (EI, 70 eV)
55 233 (M^{++} ; 86%); 203 (10%); 189 (40%); 153 (20%); 124 (40%); 77 (35%).

[0104] The product obtained BXT 51103 is of the formula below :



G/ The compound BXT 51104 of the invention is obtained, according to a method very similar to that of Example 9 from said 4,4-dimethylthieno[2,3-e]isoseleazine BXT 51103 derivative as a yellow oil.

(Beware, the product dissolved in organic solvents, such as, for example dichloromethane, slowly oxidises to give the starting material BXT 51103.)


Yield: 95%

Physical characteristics:

- * NMR ^1H : (CDCl_3)
1.2 ppm (bs, 2H); 1.37 ppm (s, 6H); 2.93 ppm (s, 2H); 7.06 ppm (d, 1H, $J = 5.3$ Hz); 7.13 ppm (d, 1H, $J = 5.3$ Hz).
- * NMR ^{13}C : (CDCl_3)
27.74 ppm; 41.35 ppm; 53.44 ppm; 119.40 ppm; 122.51 ppm; 136.86 ppm; 151.71 ppm.
- * NMR ^{77}Se : (CH_3SeCH_3)
446.6 ppm.
- * MS: (CI, isobutane)
469 (MH^+ ; 40%); 292 (20%); 276 (30%); 250 (30%); 234 (100%); 154 (30%).

Example 11: Preparation of S-(N-acetyl-L-cysteinyl)-[2-[2'-(1'-amino-2'-methyl)-propyl]-phenyl]-selenide:

[0105]



[0106] The 4,4-dimethylbenzisoseleazine derivative BXT 51072 (226 mg; 1 mmole) (see Example 8 of WO-A-95/27706) is dissolved in anhydrous methanol (5 ml). N-acetylcysteine (163 mg; 1 mmol) is added. After a few seconds, a yellow precipitate is formed and stirring is continued for 20 min. The suspension obtained is filtered and this precipitate is washed with methanol (10 ml). The desired compound is obtained as very fine yellow crystals.


Yield: 77%

Physical characteristics:

- * m.p.: 155°C (dec.)
- * NMR ^1H : (CD_3OD)
1.03 ppm (s, 6H); 1.86 ppm (s, 3H); 3.04 ppm (dd, 1H, $J = 14 - 9.5$ Hz); 3.20 ppm (m, partially superimposed with the peak of CHD_2OD); 3.65 ppm (d, 1H, $J = 13$ Hz); 3.91 ppm (d, 1H, $J = 13$ Hz); 4.19 ppm (dd, 1H, $J = 9.5 - 3.5$ Hz); 7.22-7.44 ppm (m, 3H); 8.11 ppm (m, 1H).
- * NMR ^{13}C : (CD_3OD)
22.62 ppm; 28.52 ppm; 28.86 ppm; 40.60 ppm; 40.73 ppm; 47.32 ppm; 56.20 ppm; 129.39 ppm; 130.01 ppm; 130.30 ppm; 132.72 ppm; 135.14 ppm; 143.52 ppm; 173.42 ppm; 177.57 ppm.
- * MS: (FAB; glycerol)
391 (MH^+ ; 96%); 228 (100%); 185 (100%); 150 (40%); 457 (20%).

Example 12: Preparation of S-glutathionyl-[2-[2'-(1'-amino-2'-methyl)-propyl]-phenyl]-selenide: BXT 51113

[0107]



[0108] A solution of reduced glutathione (154 mg; 0.5 mmol) in 25 ml of aqueous ethanol (7/3 v/v) is added to a solution of 4,4-dimethylbenzisoselenazine BXT 51072 (226 mg; 1 mmol) (see Example 8 of WO-A-95/27706) at ambient temperature. Stirring is kept up for 5 min.; then the solvent is evaporated under reduced pressure. The residue is taken up into 20 ml of aqueous methanol (1/2, v/v) is filtered. The filtrate is evaporated under reduced pressure, then the residue is taken up a second time in aqueous methanol, filtered, and the solvent is evaporated under reduced pressure. The desired product is obtained first as a very viscous oil, which crystallises slowly giving pale yellow crystals.

Yield: 94%

Physical characteristics:

* m.p.°C: 175°C (dec.)

* NMR ¹H: (CD₃OD/D₂O, 1/1):

1.43 ppm (s; 6H); 1.93 ppm (m; 2H); 2.32 ppm (m; 2H); 2.99 ppm (dd; 1H, J = 12 - 10 Hz); 3.25 ppm (dd; 1H, J = 13 - 4 Hz); 3.50 ppm (m; 4H); 3.68 ppm (d; 1H, J = 13 Hz); 4.70 ppm (dd; 1H, J = 10 - 4 Hz); 7.14 - 7.57 ppm (m; 3H); 7.94 ppm (m; 1H).

* NMR ¹H: (DCI/D₂O, 1N):


1.25 ppm (s; 3H); 1.27 ppm (s; 3H); 1.83 ppm (m; 2H); 2.20 ppm (m; 2H); 2.87 ppm (dd; 1H, J = 14 - 9.5 Hz); 3.08 ppm (dd; 1H, J = 14 - 4.5 Hz); 3.27 ppm (d; 1H, J = 14 Hz); 3.41 ppm (d; 1H, J = 14 Hz); 3.65 ppm (s; 2H); 3.79 ppm (t; 1H, J = 6.5 Hz); 4.19 ppm (dd; 1H, J = 9.5 - 4.5 Hz); 7.0 - 7.2 ppm (m; 3H); 7.76 ppm (m; 1H).

* MS: (FAB, glycerol)

551 (4%); 533 (M-H⁺; 5%); 459 (10%); 306 (50%); 275 (100%).

III/ Examples of synthesis of compounds of general formula II:Series wherein R₅ = hydrogen:**Example 13: Preparation of bis-4(5)-[2-phenyl-4(5)-seleno-1H-imidazole]: BXT 51038**

[0109]

**A/ Preparation of 2-phenyl-1-[2'-(trimethylsilyl)ethoxymethyl]-imidazole:**

[0110] This derivative is prepared from 2-phenyl-1H-imidazole (1.0 g; 6.9mmoles) and (trimethylsilyl)ethoxymethyl chloride (1.33 g; 7.6 mmoles) according to method (1), described in the literature by T.P. Demuth, Jr, et al. (J. Org. Chem., 1992, 57, 2963-65). The crude reaction product (2.8 g) is purified by chromatography on a silica column (eluent: cyclohexane-ethyl acetate; 6/4). The desired product is obtained as a colourless oil (1.5 g).

Yield: 80%

Physical characteristics:

* NMR ¹H: (CDCl₃)

-0.05 ppm (s, 9H); 0.87 ppm (t, 2H, J=10.50 Hz); 3.52 ppm (t, 2H, J=10.50 Hz); 5.26 ppm (s, 2H); 7.10 ppm (d, 1H, J=1.3 Hz); 7.12 ppm (d, 1H, J=1.3 Hz); 7.42 ppm (m, 3H); 7.75 ppm (m, 2H).

* NMR ¹³C: (CDCl₃)

-1.29 ppm(O); 18.0 ppm (E); 66.93 ppm (E); 75.99 ppm (E); 121.96 ppm(O); 128.27 ppm(O); 129.23 ppm(O); 129.56 ppm(O); 129.89 ppm (O); 131.46 ppm (O); 148.91 ppm (E).

* MS: (EI; 70eV)

274 (M⁺; 32); 216 (55); 158 (38); 73 (100).

B/ Preparation of bis-5-[2-phenyl-1-[2'-(trimethylsilyl)ethoxymethyl]-5-seleno-imidazole]:

[0111] A solution of selenium I chloride (0.114 g; 0.51 mmoles) in anhydrous dichloromethane (1 ml) is added dropwise, at 35-40°C and with stirring, to a solution of the preceding derivative (0.140 g; 0.51 mmoles) in the same solvent (1 ml). After a few minutes at this temperature, the red reaction mixture becomes bright yellow-orange. Then the mixture is allowed to attain ambient temperature. After one night, the yellow solution is neutralised with the aid of a saturated sodium hydrogen carbonate solution, then extracted with ethyl ether. The organic phases are combined, washed with a saturated sodium chloride solution, then dried over magnesium sulphate. After evaporation of the solvent under reduced pressure, the residual product is purified by chromatography on a silica column (eluent: cyclohexane-ethyl acetate; 7/3). The desired product is obtained as a bright yellow paste (0.210 g).

Yield: 58%

Physical characteristics:

* NMR ¹H: (CDCl₃)

-0.05 ppm (s, 9H); 0.88 ppm (t, 2H, J=10.50 Hz); 3.53 ppm (t, 2H, J=10.50 Hz); 5.25 ppm (s, 2H); 7.29 ppm (s, 1H); 7.46 ppm (m, 3H); 7.80 ppm (m, 2H).

* NMR ¹³C: (CDCl₃)

-1.40 ppm; 18.19 ppm; 66.70 ppm; 73.77 ppm; 118.26 ppm; 129.19 ppm; 129.56 ppm; 130.13 ppm; 130.65 ppm; 140.39 ppm; 153.59 ppm.

* MS: (CI; isobutane)

707 (MH⁺; 37); 355 (50); 275 (100).

C/ Preparation of bis-4(5)-[2-phenyl-4(5)-seleno-1H-imidazole]:

[0112] The preceding selenide (0.528 g; 0.75 mmoles), in solution in 30ml of a mixture of aqueous hydrofluoric acid and acetonitrile (5/95), is refluxed (=80°C) for 22h. After being cooled, the reaction mixture is neutralised with the aid of sodium hydrogen carbonate. The aqueous phase is extracted 4 times with ethyl acetate (v/v). The organic phases are combined, washed with a saturated sodium chloride solution, then dried over magnesium sulphate. After evaporation of the solvent under reduced pressure, the residual product is purified by chromatography on a silica column (eluent: cyclohexane-ethyl acetate; 6/4). The desired product is obtained as a yellow-orange solid (0.208 g).

Yield: 62%

Physical characteristics:


* m.p.°C: 219-219.5°C (hexane/ethyl acetate : 1/1)

* NMR ¹H: (CD₃OD)

7.24 ppm (s, 1H); 7.43 ppm (m, 2H); 7.80 ppm (m, 2H); 7.83 ppm (m, 2H); 7.85 ppm (m, 2H); 7.87 ppm (m, 2H); 7.89 ppm (m, 2H); 7.91 ppm (m, 2H); 7.93 ppm (m, 2H); 7.95 ppm (m, 2H); 7.97 ppm (m, 2H); 7.99 ppm (m, 2H); 8.01 ppm (m, 2H); 8.03 ppm (m, 2H); 8.05 ppm (m, 2H); 8.07 ppm (m, 2H); 8.09 ppm (m, 2H); 8.11 ppm (m, 2H); 8.13 ppm (m, 2H); 8.15 ppm (m, 2H); 8.17 ppm (m, 2H); 8.19 ppm (m, 2H); 8.21 ppm (m, 2H); 8.23 ppm (m, 2H); 8.25 ppm (m, 2H); 8.27 ppm (m, 2H); 8.29 ppm (m, 2H); 8.31 ppm (m, 2H); 8.33 ppm (m, 2H); 8.35 ppm (m, 2H); 8.37 ppm (m, 2H); 8.39 ppm (m, 2H); 8.41 ppm (m, 2H); 8.43 ppm (m, 2H); 8.45 ppm (m, 2H); 8.47 ppm (m, 2H); 8.49 ppm (m, 2H); 8.51 ppm (m, 2H); 8.53 ppm (m, 2H); 8.55 ppm (m, 2H); 8.57 ppm (m, 2H); 8.59 ppm (m, 2H); 8.61 ppm (m, 2H); 8.63 ppm (m, 2H); 8.65 ppm (m, 2H); 8.67 ppm (m, 2H); 8.69 ppm (m, 2H); 8.71 ppm (m, 2H); 8.73 ppm (m, 2H); 8.75 ppm (m, 2H); 8.77 ppm (m, 2H); 8.79 ppm (m, 2H); 8.81 ppm (m, 2H); 8.83 ppm (m, 2H); 8.85 ppm (m, 2H); 8.87 ppm (m, 2H); 8.89 ppm (m, 2H); 8.91 ppm (m, 2H); 8.93 ppm (m, 2H); 8.95 ppm (m, 2H); 8.97 ppm (m, 2H); 8.99 ppm (m, 2H); 9.01 ppm (m, 2H); 9.03 ppm (m, 2H); 9.05 ppm (m, 2H); 9.07 ppm (m, 2H); 9.09 ppm (m, 2H); 9.11 ppm (m, 2H); 9.13 ppm (m, 2H); 9.15 ppm (m, 2H); 9.17 ppm (m, 2H); 9.19 ppm (m, 2H); 9.21 ppm (m, 2H); 9.23 ppm (m, 2H); 9.25 ppm (m, 2H); 9.27 ppm (m, 2H); 9.29 ppm (m, 2H); 9.31 ppm (m, 2H); 9.33 ppm (m, 2H); 9.35 ppm (m, 2H); 9.37 ppm (m, 2H); 9.39 ppm (m, 2H); 9.41 ppm (m, 2H); 9.43 ppm (m, 2H); 9.45 ppm (m, 2H); 9.47 ppm (m, 2H); 9.49 ppm (m, 2H); 9.51 ppm (m, 2H); 9.53 ppm (m, 2H); 9.55 ppm (m, 2H); 9.57 ppm (m, 2H); 9.59 ppm (m, 2H); 9.61 ppm (m, 2H); 9.63 ppm (m, 2H); 9.65 ppm (m, 2H); 9.67 ppm (m, 2H); 9.69 ppm (m, 2H); 9.71 ppm (m, 2H); 9.73 ppm (m, 2H); 9.75 ppm (m, 2H); 9.77 ppm (m, 2H); 9.79 ppm (m, 2H); 9.81 ppm (m, 2H); 9.83 ppm (m, 2H); 9.85 ppm (m, 2H); 9.87 ppm (m, 2H); 9.89 ppm (m, 2H); 9.91 ppm (m, 2H); 9.93 ppm (m, 2H); 9.95 ppm (m, 2H); 9.97 ppm (m, 2H); 9.99 ppm (m, 2H); 10.01 ppm (m, 2H); 10.03 ppm (m, 2H); 10.05 ppm (m, 2H); 10.07 ppm (m, 2H); 10.09 ppm (m, 2H); 10.11 ppm (m, 2H); 10.13 ppm (m, 2H); 10.15 ppm (m, 2H); 10.17 ppm (m, 2H); 10.19 ppm (m, 2H); 10.21 ppm (m, 2H); 10.23 ppm (m, 2H); 10.25 ppm (m, 2H); 10.27 ppm (m, 2H); 10.29 ppm (m, 2H); 10.31 ppm (m, 2H); 10.33 ppm (m, 2H); 10.35 ppm (m, 2H); 10.37 ppm (m, 2H); 10.39 ppm (m, 2H); 10.41 ppm (m, 2H); 10.43 ppm (m, 2H); 10.45 ppm (m, 2H); 10.47 ppm (m, 2H); 10.49 ppm (m, 2H); 10.51 ppm (m, 2H); 10.53 ppm (m, 2H); 10.55 ppm (m, 2H); 10.57 ppm (m, 2H); 10.59 ppm (m, 2H); 10.61 ppm (m, 2H); 10.63 ppm (m, 2H); 10.65 ppm (m, 2H); 10.67 ppm (m, 2H); 10.69 ppm (m, 2H); 10.71 ppm (m, 2H); 10.73 ppm (m, 2H); 10.75 ppm (m, 2H); 10.77 ppm (m, 2H); 10.79 ppm (m, 2H); 10.81 ppm (m, 2H); 10.83 ppm (m, 2H); 10.85 ppm (m, 2H); 10.87 ppm (m, 2H); 10.89 ppm (m, 2H); 10.91 ppm (m, 2H); 10.93 ppm (m, 2H); 10.95 ppm (m, 2H); 10.97 ppm (m, 2H); 10.99 ppm (m, 2H); 11.01 ppm (m, 2H); 11.03 ppm (m, 2H); 11.05 ppm (m, 2H); 11.07 ppm (m, 2H); 11.09 ppm (m, 2H); 11.11 ppm (m, 2H); 11.13 ppm (m, 2H); 11.15 ppm (m, 2H); 11.17 ppm (m, 2H); 11.19 ppm (m, 2H); 11.21 ppm (m, 2H); 11.23 ppm (m, 2H); 11.25 ppm (m, 2H); 11.27 ppm (m, 2H); 11.29 ppm (m, 2H); 11.31 ppm (m, 2H); 11.33 ppm (m, 2H); 11.35 ppm (m, 2H); 11.37 ppm (m, 2H); 11.39 ppm (m, 2H); 11.41 ppm (m, 2H); 11.43 ppm (m, 2H); 11.45 ppm (m, 2H); 11.47 ppm (m, 2H); 11.49 ppm (m, 2H); 11.51 ppm (m, 2H); 11.53 ppm (m, 2H); 11.55 ppm (m, 2H); 11.57 ppm (m, 2H); 11.59 ppm (m, 2H); 11.61 ppm (m, 2H); 11.63 ppm (m, 2H); 11.65 ppm (m, 2H); 11.67 ppm (m, 2H); 11.69 ppm (m, 2H); 11.71 ppm (m, 2H); 11.73 ppm (m, 2H); 11.75 ppm (m, 2H); 11.77 ppm (m, 2H); 11.79 ppm (m, 2H); 11.81 ppm (m, 2H); 11.83 ppm (m, 2H); 11.85 ppm (m, 2H); 11.87 ppm (m, 2H); 11.89 ppm (m, 2H); 11.91 ppm (m, 2H); 11.93 ppm (m, 2H); 11.95 ppm (m, 2H); 11.97 ppm (m, 2H); 11.99 ppm (m, 2H); 12.01 ppm (m, 2H); 12.03 ppm (m, 2H); 12.05 ppm (m, 2H); 12.07 ppm (m, 2H); 12.09 ppm (m, 2H); 12.11 ppm (m, 2H); 12.13 ppm (m, 2H); 12.15 ppm (m, 2H); 12.17 ppm (m, 2H); 12.19 ppm (m, 2H); 12.21 ppm (m, 2H); 12.23 ppm (m, 2H); 12.25 ppm (m, 2H); 12.27 ppm (m, 2H); 12.29 ppm (m, 2H); 12.31 ppm (m, 2H); 12.33 ppm (m, 2H); 12.35 ppm (m, 2H); 12.37 ppm (m, 2H); 12.39 ppm (m, 2H); 12.41 ppm (m, 2H); 12.43 ppm (m, 2H); 12.45 ppm (m, 2H); 12.47 ppm (m, 2H); 12.49 ppm (m, 2H); 12.51 ppm (m, 2H); 12.53 ppm (m, 2H); 12.55 ppm (m, 2H); 12.57 ppm (m, 2H); 12.59 ppm (m, 2H); 12.61 ppm (m, 2H); 12.63 ppm (m, 2H); 12.65 ppm (m, 2H); 12.67 ppm (m, 2H); 12.69 ppm (m, 2H); 12.71 ppm (m, 2H); 12.73 ppm (m, 2H); 12.75 ppm (m, 2H); 12.77 ppm (m, 2H); 12.79 ppm (m, 2H); 12.81 ppm (m, 2H); 12.83 ppm (m, 2H); 12.85 ppm (m, 2H); 12.87 ppm (m, 2H); 12.89 ppm (m, 2H); 12.91 ppm (m, 2H); 12.93 ppm (m, 2H); 12.95 ppm (m, 2H); 12.97 ppm (m, 2H); 12.99 ppm (m, 2H); 13.01 ppm (m, 2H); 13.03 ppm (m, 2H); 13.05 ppm (m, 2H); 13.07 ppm (m, 2H); 13.09 ppm (m, 2H); 13.11 ppm (m, 2H); 13.13 ppm (m, 2H); 13.15 ppm (m, 2H); 13.17 ppm (m, 2H); 13.19 ppm (m, 2H); 13.21 ppm (m, 2H); 13.23 ppm (m, 2H); 13.25 ppm (m, 2H); 13.27 ppm (m, 2H); 13.29 ppm (m, 2H); 13.31 ppm (m, 2H); 13.33 ppm (m, 2H); 13.35 ppm (m, 2H); 13.37 ppm (m, 2H); 13.39 ppm (m, 2H); 13.41 ppm (m, 2H); 13.43 ppm (m, 2H); 13.45 ppm (m, 2H); 13.47 ppm (m, 2H); 13.49 ppm (m, 2H); 13.51 ppm (m, 2H); 13.53 ppm (m, 2H); 13.55 ppm (m, 2H); 13.57 ppm (m, 2H); 13.59 ppm (m, 2H); 13.61 ppm (m, 2H); 13.63 ppm (m, 2H); 13.65 ppm (m, 2H); 13.67 ppm (m, 2H); 13.69 ppm (m, 2H); 13.71 ppm (m, 2H); 13.73 ppm (m, 2H); 13.75 ppm (m, 2H); 13.77 ppm (m, 2H); 13.79 ppm (m, 2H); 13.81 ppm (m, 2H); 13.83 ppm (m, 2H); 13.85 ppm (m, 2H); 13.87 ppm (m, 2H); 13.89 ppm (m, 2H); 13.91 ppm (m, 2H); 13.93 ppm (m, 2H); 13.95 ppm (m, 2H); 13.97 ppm (m, 2H); 13.99 ppm (m, 2H); 14.01 ppm (m, 2H); 14.03 ppm (m, 2H); 14.05 ppm (m, 2H); 14.07 ppm (m, 2H); 14.09 ppm (m, 2H); 14.11 ppm (m, 2H); 14.13 ppm (m, 2H); 14.15 ppm (m, 2H); 14.17 ppm (m, 2H); 14.19 ppm (m, 2H); 14.21 ppm (m, 2H); 14.23 ppm (m, 2H); 14.25 ppm (m, 2H); 14.27 ppm (m, 2H); 14.29 ppm (m, 2H); 14.31 ppm (m, 2H); 14.33 ppm (m, 2H); 14.35 ppm (m, 2H); 14.37 ppm (m, 2H); 14.39 ppm (m, 2H); 14.41 ppm (m, 2H); 14.43 ppm (m, 2H); 14.45 ppm (m, 2H); 14.47 ppm (m, 2H); 14.49 ppm (m, 2H); 14.51 ppm (m, 2H); 14.53 ppm (m, 2H); 14.55 ppm (m, 2H); 14.57 ppm (m, 2H); 14.59 ppm (m, 2H); 14.61 ppm (m, 2H); 14.63 ppm (m, 2H); 14.65 ppm (m, 2H); 14.67 ppm (m, 2H); 14.69 ppm (m, 2H); 14.71 ppm (m, 2H); 14.73 ppm (m, 2H); 14.75 ppm (m, 2H); 14.77 ppm (m, 2H); 14.79 ppm (m, 2H); 14.81 ppm (m, 2H); 14.83 ppm (m, 2H); 14.85 ppm (m, 2H); 14.87 ppm (m, 2H); 14.89 ppm (m, 2H); 14.91 ppm (m, 2H); 14.93 ppm (m, 2H); 14.95 ppm (m, 2H); 14.97 ppm (m, 2H); 14.99 ppm (m, 2H); 15.01 ppm (m, 2H); 15.03 ppm (m, 2H); 15.05 ppm (m, 2H); 15.07 ppm (m, 2H); 15.09 ppm (m, 2H); 15.11 ppm (m, 2H); 15.13 ppm (m, 2H); 15.15 ppm (m, 2H); 15.17 ppm (m, 2H); 15.19 ppm (m, 2H); 15.21 ppm (m, 2H); 15.23 ppm (m, 2H); 15.25 ppm (m, 2H); 15.27 ppm (m, 2H); 15.29 ppm (m, 2H); 15.31 ppm (m, 2H); 15.33 ppm (m, 2H); 15.35 ppm (m, 2H); 15.37 ppm (m, 2H); 15.39 ppm (m, 2H); 15.41 ppm (m, 2H); 15.43 ppm (m, 2H); 15.45 ppm (m, 2H); 15.47 ppm (m, 2H); 15.49 ppm (m, 2H); 15.51 ppm (m, 2H); 15.53 ppm (m, 2H); 15.55 ppm (m, 2H); 15.57 ppm (m, 2H); 15.59 ppm (m, 2H); 15.61 ppm (m, 2H); 15.63 ppm (m, 2H); 15.65 ppm (m, 2H); 15.67 ppm (m, 2H); 15.69 ppm (m, 2H); 15.71 ppm (m, 2H); 15.73 ppm (m, 2H); 15.75 ppm (m, 2H); 15.77 ppm (m, 2H); 15.79 ppm (m, 2H); 15.81 ppm (m, 2H); 15.83 ppm (m, 2H); 15.85 ppm (m, 2H); 15.87 ppm (m, 2H); 15.89 ppm (m, 2H); 15.91 ppm (m, 2H); 15.93 ppm (m, 2H); 15.95 ppm (m, 2H); 15.97 ppm (m, 2H); 15.99 ppm (m, 2H); 16.01 ppm (m, 2H); 16.03 ppm (m, 2H); 16.05 ppm (m, 2H); 16.07 ppm (m, 2H); 16.09 ppm (m, 2H); 16.11 ppm (m, 2H); 16.13 ppm (m, 2H); 16.15 ppm (m, 2H); 16.17 ppm (m, 2H); 16.19 ppm (m, 2H); 16.21 ppm (m, 2H); 16.23 ppm (m, 2H); 16.25 ppm (m, 2H); 16.27 ppm (m, 2H); 16.29 ppm (m, 2H); 16.31 ppm (m, 2H); 16.33 ppm (m, 2H); 16.35 ppm (m, 2H); 16.37 ppm (m, 2H); 16.39 ppm (m, 2H); 16.41 ppm (m, 2H); 16.43 ppm (m, 2H); 16.45 ppm (m, 2H); 16.47 ppm (m, 2H); 16.49 ppm (m, 2H); 16.51 ppm (m, 2H); 16.53 ppm (m, 2H); 16.55 ppm (m, 2H); 16.57 ppm (m, 2H); 16.59 ppm (m, 2H); 16.61 ppm (m, 2H); 16.63 ppm (m, 2H); 16.65 ppm (m, 2H); 16.67 ppm (m, 2H); 16.69 ppm (m, 2H); 16.71 ppm (m, 2H); 16.73 ppm (m, 2H); 16.75 ppm (m, 2H); 16.77 ppm (m, 2H); 16.79 ppm (m, 2H); 16.81 ppm (m, 2H); 16.83 ppm (m, 2H); 16.85 ppm (m, 2H); 16.87 ppm (m, 2H); 16.89 ppm (m, 2H); 16.91 ppm (m, 2H); 16.93 ppm (m, 2H); 16.95 ppm (m, 2H); 16.97 ppm (m, 2H); 16.99 ppm (m, 2H); 17.01 ppm (m, 2H); 17.03 ppm (m, 2H); 17.05 ppm (m, 2H); 17.07 ppm (m, 2H); 17.09 ppm (m, 2H); 17.11 ppm (m, 2H); 17.13 ppm (m, 2H); 17.15 ppm (m, 2H); 17.17 ppm (m, 2H); 17.19 ppm (m, 2H); 17.21 ppm (m, 2H); 17.23 ppm (m, 2H); 17.25 ppm (m, 2H); 17.27 ppm (m, 2H); 17.29 ppm (m, 2H); 17.31 ppm (m, 2H); 17.33 ppm (m, 2H); 17.35 ppm (m, 2H); 17.37 ppm (m, 2H); 17.39 ppm (m, 2H); 17.41 ppm (m, 2H); 17.43 ppm (m, 2H); 17.45 ppm (m, 2H); 17.47 ppm (m, 2H); 17.49 ppm (m, 2H); 17.51 ppm (m, 2H); 17.53 ppm (m, 2H); 17.55 ppm (m, 2H); 17.57 ppm (m, 2H); 17.59 ppm (m, 2H); 17.61 ppm (m, 2H); 17.63 ppm (m, 2H); 17.65 ppm (m, 2H); 17.67 ppm (m, 2H); 17.69 ppm (m, 2H); 17.71 ppm (m, 2H); 17.73 ppm (m, 2H); 17.75 ppm (m, 2H); 17.77 ppm (m, 2H); 17.79 ppm (m, 2H); 17.81 ppm (m, 2H); 17.83 ppm (m, 2H); 17.85 ppm (m, 2H); 17.87 ppm (m, 2H); 17.89 ppm (m, 2H); 17.91 ppm (m, 2H); 17.93 ppm (m, 2H); 17.95 ppm (m, 2H); 17.97 ppm (m, 2H); 17.99 ppm (m, 2H); 18.01 ppm (m, 2H); 18.03 ppm (m, 2H); 18.05 ppm (m, 2H); 18.07 ppm (m, 2H); 18.09 ppm (m, 2H); 18.11 ppm (m, 2H); 18.13 ppm (m, 2H); 18.15 ppm (m, 2H); 18.17 ppm (m, 2H); 18.19 ppm (m, 2H); 18.21 ppm (m, 2H); 18.23 ppm (m, 2H); 18.25 ppm (m, 2H); 18.27 ppm (m, 2H); 18.29 ppm (m, 2H); 18.31 ppm (m, 2H); 18.33 ppm (m, 2H); 18.35 ppm (m, 2H); 18.37 ppm (m, 2H); 18.39 ppm (m, 2H); 18.41 ppm (m, 2H); 18.43 ppm (m, 2H); 18.45 ppm (m, 2H); 18.47 ppm (m, 2H); 18.49 ppm (m, 2H); 18.51 ppm (m, 2H); 18.53 ppm (m, 2H); 18.55 ppm (m, 2H); 18.57 ppm (m, 2H); 18.59 ppm (m, 2H); 18.61 ppm (m, 2H); 18.63 ppm (m, 2H); 18.65 ppm (m, 2H); 18.67 ppm (m, 2H); 18.69 ppm (m, 2H); 18.71 ppm (m, 2H); 18.73 ppm (m, 2H); 18.75 ppm (m, 2H); 18.77 ppm (m, 2H); 18.79 ppm (m, 2H); 18.81 ppm (m, 2H); 18.83 ppm (m, 2H); 18.85 ppm (m, 2H); 18.87 ppm (m, 2H); 18.89 ppm (m, 2H); 18.91 ppm (m, 2H); 18.93 ppm (m, 2H); 18.95 ppm (m, 2H); 18.97 ppm (m, 2H); 18.99 ppm (m, 2H); 19.01 ppm (m, 2H); 19.03 ppm (m, 2H); 19.05 ppm (m, 2H); 19.07 ppm (m, 2H); 19.09 ppm (m, 2H); 19.11 ppm (m, 2H); 19.13 ppm (m, 2H); 19.15 ppm (m, 2H); 19.17 ppm (m, 2H); 19.19 ppm (m, 2H); 19.21 ppm (m, 2H); 19.23 ppm (m, 2H); 19.25 ppm (m, 2H); 19.27 ppm (m, 2H); 19.29 ppm (m, 2H); 19.31 ppm (m, 2H); 19.33 ppm (m, 2H); 19.35 ppm (m, 2H); 19.37 ppm (m, 2H); 19.39 ppm (m, 2H); 19.41 ppm (m, 2H); 19.43 ppm (m, 2H); 19.45 ppm (m, 2H); 19.47 ppm (m, 2H); 19.49 ppm (m, 2H); 19.51 ppm (m, 2H); 19.53 ppm (m, 2H); 19.55 ppm (m, 2H); 19.57 ppm (m, 2H); 19.59 ppm (m, 2H); 19.61 ppm (m, 2H); 19.63 ppm (m, 2H); 19.65 ppm (m, 2H); 19.67 ppm (m, 2H); 19.69 ppm (m, 2H); 19.71 ppm (m, 2H); 19.73 ppm (m, 2H); 19.75 ppm (m, 2H); 19.77 ppm (m, 2H); 19.79 ppm (m, 2H); 19.81 ppm (m, 2H); 19.83 ppm (m, 2H); 19.85 ppm (m, 2H); 19.87 ppm (m, 2H); 19.89 ppm (m, 2H); 19.91 ppm (m, 2H); 19.93 ppm (m, 2H); 19.95 ppm (m, 2H); 19.97 ppm (m, 2H); 19.99 ppm (m, 2H); 20.01 ppm (m, 2H); 20.03 ppm (m, 2H); 20.05 ppm (m, 2H); 20.07 ppm (m, 2H); 20.09 ppm (m, 2H); 20.11 ppm (m, 2H); 20.13 ppm (m, 2H); 20.15 ppm (m, 2H); 20.17 ppm (m, 2H); 20.19 ppm (m, 2H); 20.21 ppm (m, 2H); 20.23 ppm (m, 2H); 20.25 ppm (m, 2H); 20.27 ppm (m, 2H); 20.29 ppm (m, 2H); 20.31 ppm (m, 2H); 20.33 ppm (m, 2H); 20.35 ppm (m, 2H); 20.37 ppm (m, 2H); 20.39 ppm (m, 2H); 20.41 ppm (m, 2H); 20.43 ppm (m, 2H); 20.45 ppm (m, 2H); 20.47 ppm (m, 2H); 20.49 ppm (m, 2H); 20.51 ppm (m, 2H); 20.53 ppm (m, 2H); 20.55 ppm (m, 2H); 20.57 ppm (m, 2H); 20.59 ppm (m, 2H); 20.61 ppm (m, 2H); 20.63 ppm (m, 2H); 20.65 ppm (m, 2H); 20.67 ppm (m, 2H); 20.69 ppm (m, 2H); 20.71 ppm (m, 2H); 20.73 ppm (m, 2H); 20.75 ppm (m, 2H); 20.77 ppm (m, 2H); 20.79 ppm (m, 2H); 20.81 ppm (m, 2H); 20.83 ppm (m, 2H); 20.85 ppm (m, 2H); 20.87 ppm (m, 2H); 20.89 ppm (m, 2H); 20.91 ppm (m, 2H); 20.93 ppm (m, 2H); 20.95 ppm (m, 2H); 20.97 ppm (m, 2H); 20.99 ppm (m, 2H); 21.01 ppm (m, 2H); 21.03 ppm (m, 2H); 21.05 ppm (m, 2H); 21.07 ppm (m, 2H); 21.09 ppm (m, 2H); 21.11 ppm (m, 2H); 21.13 ppm (m, 2H); 21.15 ppm (m, 2H); 21.17 ppm (m, 2H); 21.19 ppm (m, 2H); 21.21 ppm (m, 2H); 21.23 ppm (m, 2H); 21.25 ppm (m, 2H); 21.27 ppm (m, 2H); 21.29 ppm (m, 2H); 21.31 ppm (m, 2H); 21.33 ppm (m, 2H); 21.35 ppm (m, 2H); 21.37 ppm (m, 2H); 21.39 ppm (m, 2H); 21.41 ppm (m, 2H); 21.43 ppm (m, 2H); 21.45 ppm (m, 2H); 21.47 ppm (m, 2H); 21.49 ppm (m, 2H); 21.51 ppm (m, 2H); 21.53 ppm (m, 2H); 21.55 ppm (m, 2H); 21.57 ppm (m, 2H); 21.59 ppm (m, 2H); 21.61 ppm (m, 2H); 21.63 ppm (m, 2H); 21.65 ppm (m, 2H); 21.67 ppm (m, 2H); 21.69 ppm (m, 2H); 21.71 ppm (m, 2H); 21.73 ppm (m, 2H); 21.75 ppm (m, 2H); 21.77 ppm (m, 2H); 21.79 ppm (m, 2H); 21.81 ppm (m, 2H); 21.83 ppm (m, 2H); 21.85 ppm (m, 2H); 21.87 ppm (m, 2H); 21.89 ppm (m, 2H); 21.91 ppm (m, 2H); 21.93 ppm (m, 2H); 21.95 ppm (m, 2H); 21.97 ppm (m, 2H); 21.99 ppm (m, 2H); 22.01 ppm (m, 2H); 22.03 ppm (m, 2H); 22.05 ppm (m, 2H); 22.07 ppm (m, 2H); 22.09 ppm (m, 2H); 22.11 ppm (m, 2H); 22.13 ppm (m, 2H); 22.15 ppm (m, 2H); 22.17 ppm (m, 2H); 22.19 ppm (m, 2H); 22.21 ppm (m, 2H); 22.23 ppm (m, 2H); 22.25 ppm (m, 2H); 22.27 ppm (m, 2H); 22.29 ppm (m, 2H); 22.31 ppm (m, 2H); 22.33 ppm (m, 2H); 22.35 ppm (m, 2H); 22.37 ppm (m, 2H); 22.39 ppm (m, 2H); 22.41 ppm (m, 2H); 22.43 ppm (m, 2H); 22.45 ppm (m, 2H); 22.47 ppm (m, 2H); 22.49 ppm (m, 2H); 22.51 ppm (m, 2H); 22.53 ppm (m, 2H); 22.55 ppm (m, 2H); 22.57 ppm (m, 2H); 22.59 ppm (m, 2H); 22.61 ppm (m, 2H); 22.63 ppm (m, 2H); 22.65 ppm (m, 2H); 22.67 ppm (m, 2H); 22.69 ppm (m, 2H); 22.71 ppm (m, 2H); 22.73 ppm (m, 2H); 22.75 ppm (m, 2H); 22.77 ppm (m, 2H); 22.79 ppm (

Example 14: Preparation of bis-4(5)-[[2-(4'-carbomethoxy)phenyl]-4(5)-seleno-1H-imidazole]: BXT 51079

[0113]

**A/ Preparation of [2-[(4'-carbomethoxy)phenyl]-1-[2'-(trimethylsilyl)ethoxy-methyl]-imidazole:**

[0114] The preparation of the intermediate [1-(trimethylsilyl)ethoxymethyl]-imidazole-2-zinc chloride is carried out according to method (2), described in the literature by A. S. Bell, D. A. Roberts, and K. S. Ruddock (Tetrahedron Lett., 1988, 29(39), 5013-16), from 1-(trimethylsilyl)ethoxymethyl-imidazole (6.075 g; 30.8 mmoles), (prepared according to method (1) already cited in Example 13/A), and anhydrous zinc chloride (5.0 g; 36.7 mmoles). The organozinc derivative thus obtained, is used directly for the following coupling reaction and is not isolated.

[0115] The suspension of the preceding intermediate [1-(trimethylsilyl)ethoxymethyl]-imidazole-2-zinc chloride, (55 ml; 30.8 mmoles) in anhydrous THF, is treated under nitrogen, with palladium (0) tetrakis(triphenyl)phosphine (0.4g; 0.35mmoles), and methyl 4-bromobenzoate (4.6g; 21.6mmoles) in anhydrous THF (15 ml). The reaction mixture is heated one hour under reflux (~65-70°C), under nitrogen, then 8 additional hours, after addition of an excess of zinc chloride (8.8g; 61 mmoles). The reaction mixture is then treated in a similar manner to the literature (2). After evaporation of the solvent under reduced pressure, the residual product is purified by chromatography on a silica column (eluent gradient: cyclohexane-ethyl acetate 85/15; then ethyl acetate-cyclohexane 95/5; then ethyl acetate-methanol 90/10). The desired product is obtained as a white solid (5.56 g).

Yield: 90%

Physical characteristics:

* NMR ¹H: (CDCl₃)

0.002 ppm (s, 9H); 0.94 ppm (t, 2H, J=8.60 Hz); 3.61 ppm (t, 2H, J=8.60 Hz); 3.94 ppm (s, 3H); 5.30 ppm (s, 2H); 7.15 ppm (d, 1H, J=1.42 Hz); 7.18 ppm (d, 1H, J=1.42 Hz); 7.92 ppm (m symmetrical, 2H); 8.12 ppm (m symmetrical, 2H).

* NMR ¹³C: (CDCl₃)

-1.3 ppm; 17.99 ppm; 52.55 ppm; 66.95 ppm; 75.93 ppm; 122.94 ppm; 129.16 ppm; 129.62 ppm; 130.38 ppm; 130.73 ppm; 134.96 ppm; 148.017 ppm; 167.38 ppm.

* MS: (EI; 70eV)

332 (M⁺; 28); 289 (32); 274 (68); 170 (86); 73 (100).**B/ Preparation of bis-5-[[2-(4'-carbomethoxy)phenyl]-1-[2'-(trimethylsilyl)ethoxymethyl]-5-seleno-imidazole]:**

[0116] The desired product is obtained from the preceding derivative (4.6 g, 14 mmoles) and selenium I chloride (7.6 g, 33.3 mmoles) in anhydrous dichloromethane (80 ml) in following the same procedure as that described in Example 13/B (the temperature of addition of Se₂Cl₂ is 0° C in the present case). The desired product is obtained after purification by chromatography on an alumina column (eluent gradient: pure dichloromethane, then dichloromethane/ethyl acetate 90/10, then pure ethyl acetate, then ethyl acetate/ methanol 90/10), as a bright yellow solid (2.46 g).

Yield: 43 %

Physical characteristics:

* NMR ¹H: (CDCl₃)

0.005 ppm (s, 9H); 0.95 ppm (t, 2H, J=9.0 Hz); 3.61 ppm (t, 2H, J=9.0 Hz); 3.96 ppm (s, 3H); 5.32 ppm (s, 2H); 7.33 ppm (s, 1H); 7.96 ppm (m symmetrical, 2H); 8.15 ppm (m symmetrical, 2H).

* NMR ¹³C: (CDCl₃)

1.6 ppm; 18.26 ppm; 52.65 ppm; 53.75 ppm; 66.98 ppm; 73.86 ppm; 119.11 ppm; 129.39 ppm; 130.46 ppm; 131.45 ppm; 134.60 ppm; 140.569 ppm; 152.47 ppm; 167.18 ppm.

* MS: (FAB +; glycerol)

823 (MH⁺; 85); 743 (M-Se, 12); 543 (45); 375 (100).

C/ Preparation of bis-4(5)-[[2-(4'-carbomethoxy)phenyl]-4(5)-seleno-1H-imidazole]:

[0117] The preceding selenide (0.260 g ; 0.32 mmoles) is treated following the same method as that described in Example 13/C. The desired product is obtained after purification by chromatography on an alumina column (eluent gradient: ethyl acetate/methanol (0.5 %, then 1 %) as a yellow orange solid (0.108 g).

Yield: 60 %

Physical characteristics:

* NMR ¹H: (CD₃OD)

3.94 ppm (s, 3H); 7.35 ppm (bs, 1H); 7.97 ppm (m symmetrical, 2H); 8.07 (m symmetrical, 2H)

* NMR ¹³C: (CD₃OD)


52.90 ppm; 126.65 ppm; 126.88 ppm; 126.94 ppm; 131.55 ppm; 131.96 ppm; 135.16 ppm; 166.61 ppm; 168.34 ppm.

* MS: (CI; isobutane)

563 (MH⁺; 2); 483 (M-Se, 1.8); 203(100).

Example 15 : Preparation of bis-4(5)-[2-[(4'-carboxyl)phenyl]-4(5)-seleno-1H-imidazole]: BXT 51043

[0118]



[0119] The derivative BXT 51079 (0.080 g ; 0.14 mmoles), obtained in the preceding Example, is dissolved in anhydrous THF (16 ml). To this solution cooled at 0°C is added an aqueous solution (8 ml) of lithium hydroxide (0.075 g, 1.8 mmoles). The reaction is continued for 24 hours at ambient temperature. After the treatment of the crude reaction mixture, according to method (3) described in the literature by D. A. Evans et al., (Tetrahedron., 1988, 17, 5525-5540), the desired product is obtained as a yellow-green solid (0.154 g).

Yield: 90%

Physical characteristics:

* m.p.°C: ~ 230°C (dec)

* NMR ¹H: (CD₃OD)


7.28 ppm (bs, 1H); 7.92 ppm (m, 2H); 8.05(m, 2H).

* NMR ¹³C: (CD₃OD)

126.21 ppm; 126.4 ppm; 126.6 ppm; 131.31 ppm; 132.58 ppm; 137.0 ppm; 140.0 ppm; 174.89 ppm;

Example 16 : Preparation of bis-4(5)-[2-(4'-carbo(2''-(4'''-methyl-piperazin-1''-yl)ethoxyphenyl)]-4(5)-seleno-1H-imidazole] : BXT 51070

[0120]



A/ Preparation of bis-5-[[2-(4'-carboxyl)phenyl]-1-[2'-(trimethylsilyl)ethoxy-methyl]-5-seleno-imidazole]:

[0121] The desired product is obtained from the derivative prepared in Example 14/A (2.1 g, 2.56 mmoles) in solution in anhydrous THF (20 ml), treated with an aqueous solution (10 ml) of lithium hydroxide (0.260 g, 12.3 mmoles) according to method (3) already cited. After a lyophilisation step, the compound is obtained as a yellow crystalline product (2.4 g).

Yield: quantitative

Physical characteristics:

- * NMR ¹H: (CD₃OD)
-0.02 ppm (s, 9H); 0.92 ppm (t, 2H, J=7.7 Hz); 3.60 ppm (t, 2H, J=7.7 Hz); 5.42 ppm (s, 2H); 7.29 ppm (s, 1H); 7.81 ppm (m symmetrical, 2H); 8.11 ppm (m symmetrical, 2H).
- * NMR ¹³C: (CD₃OD)
1.6 ppm (O); 18.83 ppm (E); 67.80 ppm (E); 75.24 ppm (E); 120.43 ppm (E); 135.03 (O); 131.11 (O); 132.69 ppm (E); 140.04 (O); 140.89 (E); 154.42 ppm (E); 174.52 (E).
- * MS: (FAB+; glycerol)
795 (MH⁺; 10); 399 (100).

B/ Preparation of bis-5-[[2-(4'carbosuccinimidyloxy)phenyl]-1[2'-(trimethyl-silyl)ethoxymethyl]-5-seleno-imidazole].

[0122] This derivative is obtained from the preceding product (0.400 g ; 0.5 mmoles) and N-hydroxysuccinimide (0.115 g ; 1 mmole) according to method described in the literature by G. W. Anderson et al., (J. Am. Chem. Soc., 1964, 86, 1839) and W. König and R. Geiger (Chem. Ber., 1973, 106, 3626). After 72 hours at 25 °C, the reaction mixture is treated in an identical manner to that described. The solvent is evaporated under reduced pressure, and the crude product is isolated, as a bright yellow-orange solid (0.495 g).

Yield: 76 %

Physical characteristics:

- * NMR ¹H: (CDCl₃)
-0.015 ppm (s, 9H); 0.97 ppm (t, 2H, J=8.4 Hz); 2.94 ppm (bs, 4H); 3.66 ppm (t, 2H, J=8.18 Hz); 5.37 ppm (s, 2H); 7.35 ppm (s, 1H); 8.09 ppm (m symmetrical, 2H); 8.27 ppm (m symmetrical, 2H).
- * MS: (FAB+; para-nitrobenzyl alcohol)
990 (MH⁺; 14); 495 (MH⁺/2; 20); 439 (100).

C/ Preparation of bis-5-[2[4'-carbo(2''-(4'''-methyl-piperazin-1''-yl)ethoxyphenyl)]-5-seleno-1-[2'-(trimethylsilyl)ethoxymethyl]-1H-imidazole]:

[0123]

1/ Preparation of N-(2-hydroxyethyl)-N'-methyl-piperazine:

N-methyl-piperazine (10 g, 100 mmoles) and 2-chloroethanol (8.5 g, 100 mmoles) are stirred at 100 °C for 4

hours. 250 ml of acetone are added to the very viscous reaction mixture and the resulting suspension is neutralised with 15 ml of triethylamine. After filtration of the triethylammonium chloride, the solvent is evaporated under reduced pressure. The desired compound is obtained after purification by chromatography on an alumina column (eluent: ethyl acetate) as a colourless oil.

Yield: 75 %

Physical characteristics:

* NMR ¹H: (CDCl₃)

2.20 ppm(s, 3H); 2.39 ppm(m, 8H); 2.46 ppm(t, 2H, J=5.5 Hz); 3.41 ppm(bs, 1H); 3.54 ppm(t, 2H, J=5.5 Hz)

2/ - Preparation of bis-5-[2[4'-carbo(2''-(4'''-methyl-piperazin-1''-yl)-ethoxyphenyl]-5-seleno-1-[2-(trimethylsilyl)ethoxymethyl]-1H-imidazole].

N-hydroxysuccinimide ester, prepared in Example 17/B (0.270 g, 0.275 mmoles) and dissolved in a minimum of anhydrous THF (2 ml), is mixed with N-2-hydroxyethyl-N'-methylpiperazine (2.15 g, 3.5 mmoles), used here as solvent. After addition at ambient temperature of titanium tetrakisopropoxide (0.055 ml, 0.137 mmoles), the reaction mixture is heated at 80° C for 2 hours. Once the reaction has finished, the THF present in the medium is evaporated, then the excess of alcohol is distilled off under reduced pressure (T = 90-100°C/p=0.2 mmHg). The residue is finally purified by chromatography on a basic alumina column - activity I (eluent: ethyl acetate/methanol 95/5). A bright yellow product is isolated (0.113 g).

Yield: 40 %

Physical characteristics:

* NMR ¹H: (CDCl₃)

-0.041 ppm (s, 9H); 0.89 ppm (t, 2H, J=8.14 Hz); 2.31 ppm (s, 3H); 2.62 ppm(m, 8H); 2.83 ppm (t, 2H, J=5.4 Hz); 5.39 ppm (s, 2H); 7.30 ppm (bs, 1H); 7.92 ppm (m symmetrical, 2H); 8.16 ppm(m symmetrical, 2H).

* NMR ¹³C: (CD₃OD)

-1.30 ppm(O); 18.86 ppm(E); 45.90 ppm(O); 53.82 ppm(O); 55.71 ppm(E); 57.66 ppm(E); 63.64 ppm(E); 67.90 (E); 75.18 (E); 121.0 ppm(E); 130.59 ppm(O); 131.38 ppm(O); 132.99 ppm(E); 135.45 ppm(E); 140.38 ppm (O); 153.41 ppm(E); 167.51 ppm(E).

* MS: (FAB+, glycerol)

1047 (MH⁺; 3.5); 525(MH⁺/2; 80); 445 (MH⁺/2-Se, 58)

D/ Preparation of bis-4(5)-[2[4'-carbo(2''-(4'''-methyl-piperazin-1''-yl)ethoxy phenyl]-4(5)-seleno-1H-imidazole].

[0124] The preceding selenide (0.143 g, 0.137 mmoles) in solution in 14 ml of a mixture of aqueous hydrofluoric acid and acetonitrile (5/95) is refluxed (~ 80°C) for 20 hours. After being cooled, the reaction mixture is treated in the same way as that described in Example 13/C. After having evaporated the solvent under reduced pressure, the residue is then purified by chromatography on a basic alumina column - activity I (eluent gradient: ethyl acetate/methanol 95/5, then ethyl acetate/methanol 90/10). The desired product is obtained as a bright yellow paste (0.049 g).

Yield: 45 %

Physical characteristics:

* NMR ¹H: (CD₃OD)

2.26 ppm (s, 3H); 2.60 ppm(m, 8H); 2.82 ppm (t, 2H, J=5.4 Hz); 4.48 ppm (s, 2H); 7.33 ppm (bs, 1H); 7.97 ppm (m symmetrical, 2H); 8.10 ppm (m symmetrical, 2H).

* NMR ¹³C: (CD₃OD)

45.67 ppm(O); 53.90 ppm(O); 54.91 ppm(E); 56.66 ppm(E); 62.89 ppm(E); 126.26 ppm(E); 126.53 ppm(O); 130.35 ppm(O); 130.62 ppm(E); 134.15 ppm(E); 149.35 ppm(O); 166.39 ppm(E); 177.52(E).


* MS: (FAB+, glycerol)

785 (MH⁺; 4); 393(MH⁺/2; 33)

Series wherein $R_5 \neq$ hydrogen:

Example 17: Preparation of bis-5-[2-[(4'-trifluoromethyl)phenyl]-1-methyl-imidazole]: BXT 51045

[0125]



A/ Preparation of 2-[(4'-trifluoromethyl)phenyl]-1-methyl-imidazole:

[0126] The preparation of the intermediate 2-(tributylstannyl)-1-methyl-imidazole is carried out according to method described by K. C. Molloy, et al., (J. of Organomet. Chem., 1989, 365 61-73), from 1-methylimidazole (0.16 ml; 2 mmoles) and tributyltin chloride (0.56 ml; 2 mmoles). The stannyl derivative thus obtained is used directly for the following coupling reaction and is not isolated.

[0127] A solution of the preceding 2-(tributylstannyl)-1-methylimidazole (5 ml; 2 mmoles) in anhydrous THF is added, dropwise under argon, onto a suspension of dichlorobis(triphenylphosphine) palladium(II) (0.120 g; 0.17 mmoles) and 4-bromo-trifluoromethylbenzene (0.23 ml; 1.7 mmoles) in m-xylene (4 ml). The reaction mixture is refluxed ($\approx 120^\circ\text{C}$), under argon, for 20h. After having been washed with the aid of an aqueous solution of potassium fluoride, the reaction mixture is extracted 4 times with ethyl ether (v/v). The organic phases are combined, washed with a saturated sodium chloride solution, then dried over magnesium sulphate. After evaporation of the solvent under reduced pressure, the residual product is purified by chromatography on a silica column (eluent gradient: cyclohexane-ethyl acetate 30/70; then ethyl acetate 100; then ethyl acetate/methanol 95/5). The desired product is obtained as a beige solid (0.223 g). Yield: 58%

Physical characteristics:

- * m.p. $^\circ\text{C}$: 115-116 $^\circ\text{C}$ (ethyl acetate)
- * NMR ^1H : (CDCl_3)
3.79 ppm (s, 3H); 7.02 ppm (d, 1H; $J=1.15$ Hz); 7.16 ppm (d, 1H; $J=1.15$ Hz); 7.74 ppm (m symmetrical, 4H).
- * NMR ^{13}C : (CDCl_3)
34.89 ppm (O); 123.68 ppm (O); 125.99 ppm (O); 126.03 ppm (E) (q, $^2J_{\text{C-F}} = 32$ Hz); 128.02 ppm (O) (q, $^1J_{\text{C-F}} = 207$ Hz); 129.28 ppm (O); 129.54 ppm (O); 130.65 ppm (E); 134.62 ppm (E).
- * MS: (EI, 70eV)
225 (M^{++} ; 100); 156(6); 145(5); 113 (8); 84 (14); 49 (18).

B/ Preparation of bis-5-[2-[(4'-trifluoromethyl)phenyl]-1-methyl-imidazole]:

[0128] The desired product is obtained from the preceding derivative (0.198 g; 0.876 mmoles) and selenium I chloride (0.252 g; 1.1 moles) in anhydrous dichloromethane in following the same procedure as that described in Example 13/B. The desired product is obtained after purification by chromatography on a silica column (eluent gradient: ethyl acetate/cyclohexane 7/3, then pure ethyl acetate) as a yellow-orange solid (0.2123 g)

Yield: 79%


Physical characteristics:

- * m.p. $^\circ\text{C}$: 168-169 $^\circ\text{C}$ (ethyl acetate; partial sublimation)
- * NMR ^1H : (CDCl_3)
3.71 ppm (s, 3H); 7.34 ppm (s, 1H); 7.72-7.83 ppm (m, 4H).
- * NMR ^{13}C : (CDCl_3)
33.90 ppm (O); 118.81 ppm (E); 126.19 ppm (O); 126.22 ppm (E) (q, $^2J_{\text{C-F}} = 32$ Hz); 127.9 ppm (O) (q, $^1J_{\text{C-F}} = 235$ Hz); 129.56 ppm (O); 134.29 ppm (E); 140.26 ppm (O); 151.05 ppm (E).

- * NMR ^{77}Se (CH_3SeCH_3)
401.6 ppm
- * MS: (EI, 70eV)
610 (M^{+} ; 8); 450 (12); 305 (100).

Example 18: Preparation of bis-5-[2-[(4'-carbomethoxy)phenyl]-5-seleno-1-methyl-imidazole]: BXT 51047

[0129]



A/ Preparation of 2-[(4'-carbomethoxy)phenyl]-1-methyl-imidazole:

[0130] This derivative is obtained by a coupling reaction between 2-(tributylstannyl)-1-methylimidazole (whose preparation is identical to that used in Example 17/A) in solution in THF (5 ml; 2.0 mmoles) and methyl 4-bromobenzoate (0.365 g; 1.7 mmoles) in the presence of dichlorobis (triphenylphosphine) palladium (II) (0.060 g; 0.085 mmoles) according to the same method as that described in Example 18/A. The desired product is purified by chromatography on a silica column (eluent gradient: cyclohexane-ethyl acetate 30/70; then ethyl acetate 100; then ethyl acetate/methanol 95/5).

The desired product is obtained as a white solid (0.141 g).

Yield: 40%

Physical characteristics:

- * m.p. °C : 133-134 °C (ethyl acetate)
- * NMR ^1H : (CDCl_3)
3.77 ppm (s, 3H); 3.92 ppm (s, 3H); 6.99 ppm (d, 1H, $J=1.22$ Hz); 7.14 ppm (d, 1H, $J=1.22$ Hz); 7.72 ppm (m symmetrical, 2H); 8.10 ppm (m symmetrical, 2H).
- * NMR ^{13}C : (CDCl_3)
34.99 ppm(O); 52.55 ppm(O); 123.72 ppm(O); 128.95 ppm(O); 129.54 ppm(O); 130.32 ppm(O); 130.45 ppm(E); 135.4 ppm(E); 147.25 ppm(E); 167.32 ppm(E).
- * MS: (EI; 70eV)
216 (M^{+} ; 100); 185(29); 157(22)

B/ Preparation of bis-5-[2-[(4'-carbomethoxy)phenyl]-5-seleno-1-methyl-imidazole]:

[0131] A solution of selenium I chloride (0.30 g; 1.3 mmoles) in anhydrous dichloromethane (2.6 ml) is added dropwise, at ambient temperature and with stirring, to a solution of the preceding compound (0.280 g; 1.3 mmoles) in the same solvent (2.6 ml). After 17 h of reaction, the yellowish mixture is neutralised with the aid of a saturated sodium hydrogen carbonate solution, then extracted with ethyl ether. The organic phases are combined, washed with a saturated sodium chloride solution, then dried over MgSO_4 . After evaporation of the solvent under reduced pressure, the residual product is purified by chromatography on a silica column (eluent gradient: ethyl acetate/cyclohexane 80/20; then pure ethyl acetate).

[0132] The desired product is obtained as a bright yellow solid (0.204 g).

Yield: 70%


Physical characteristics:

- * m.p. °C: 196.5 °C (ethyl acetate)
- * NMR ^1H : (CDCl_3)
3.71 ppm (s, 3H); 3.96 ppm (s, 3H); 7.35 ppm (s, 1H); 7.75 ppm (m symmetrical, 2H); 8.15 ppm (m symmetrical, 2H).

- * NMR ^{13}C : (CDCl_3)
33.99 ppm (O); 52.65 ppm (O); 118.82 ppm (E); 129.14 ppm (O); 130.46 ppm (O); 131.21 ppm (E); 135.0 ppm (E); 140.28 ppm (O); 151.47 ppm (E); 167.10 ppm (E).
- * NMR ^{77}Se (D_2O)
402.3 ppm
- * MS: (CI; isobutane)
591 (MH^+ ; 9); 511 (M-Se, 4); 311 (100); 296 ($\text{MH}^+ - 2$, 45)

Example 19: Preparation of bis-5-[2-[(4'-carboxyl)-phenyl]-5-seleno-1-methyl-imidazole] : BXT 51046.

[0133]



[0134] The derivative BXT 51047 (0.025 g; 0.05 mmoles), obtained in Example 18/B, is dissolved in 0.5 ml of a mixture methanol-water (3:1). Cooled to 5°C, an aqueous solution of lithium hydroxide in the same proportions as the two preceding solvents (0.030 g; 0.5 mmoles), is added to this suspension. After a 17-hour reaction, at this temperature, the yellow suspension obtained is treated according to method (3), already cited in Example 16. The residual product, taken up into a minimum of methanol, is purified by chromatography on a silica column (eluent gradient: methanol/ethyl acetate 40/60, then methanol/ethyl acetate 80/20). The desired product is obtained as a yellow solid (0.028 g).

Yield: 92 %

Physical characteristics:

- * NMR ^1H : (CD_3OD)
3.76 ppm (s, 3H); 7.22 ppm (s, 1H); 7.67 ppm (m symmetrical, 2H); 8.11 ppm (m symmetrical, 2H)
- * NMR ^{13}C : (CD_3OD)
34.25 ppm (O); 119.7 ppm (E); 129.82 ppm (O); 131.02 ppm (O); 132.95 ppm (E); 139.58 ppm (O); 140.85 ppm (E); 153.54 ppm (E); 174.79 ppm (E).
- * NMR ^{77}Se (D_2O)
402.3 ppm

II/ Activities:

[0135] The operatory protocols described below are non-limiting examples of applications of the method according to the invention.

EXAMPLE 20 : MEASUREMENT OF THE GLUTATHIONE PEROXIDASE ACTIVITY OF COMPOUNDS OF GENERAL FORMULA I

[0136] The glutathione peroxidase activity is determined by using a 50 mM HEPES buffer, pH=7.3 (at 37°C), containing 0.2 mM DTPA, 0.144 mM NADPH, 2.2 mM reduced glutathione (GSH) and 1.1 U/ml glutathione disulphide reductase. This buffer further contains 110 U/ml catalase when hydrogen peroxide is not used as substrate.

[0137] To 1.5 ml of the buffer described above, are added 100 μl of an ethanolic stock solution of the compound tested or 100 μl of absolute ethanol (blank). Each compound is tested at a final concentration of 10 μM . The reaction medium is equilibrated for 2 minutes at 37°C. Then 50 μl of 6.6 mM tert-butyl hydroperoxide (t-BuOOH) in ultrapure water, or 3.3 mM hydrogen peroxide (H_2O_2) in ultrapure water are added. The glutathione peroxidase activity is determined at 37°C by measuring the decrease of absorbance at 340 nm for 5 minutes. Said activity or initial enzymatic rate is proportional to the slope of the variation of absorbance with time.

[0138] The catalytic activity of oxygen reduction of the compounds tested corresponds to the rate of consumption of

NADPH in the absence of hydroperoxide. When this rate is significantly greater than that of the control, the corresponding glutathione oxidase activity can be checked by the direct measurement of the kinetics of consumption of the dissolved oxygen with the aid of a Clark electrode.

[0139] The results of the glutathione peroxidase activity measurements are shown in Table 1 below. They are expressed in nmoles of hydroperoxide reduced per minute.

TABLE 1

Glutathione peroxidase activity (in nmoles of hydroperoxide reduced/min) pH = 7.3; 37°C; [GSH] = 2 mM		
	t-BuOOH	H ₂ O ₂
BXT-51079	7.9	13.9
BXT-51099	9.6	31.1
BXT-51104	16.8	44.7
BXT-51108	24.7	46.8
BXT-51110	14.3	n.d.
BXT-51111	12.9	n.d.
BXT-51125	17.7	n.d.
n.d. = non-determined		

[0140] As shown in Table 1 the compounds of general structure I described in the present invention catalyze, in the presence of glutathione GSH, the reduction of hydrogen peroxide or that of an organic hydroperoxide.

EXAMPLE 21: INHIBITION OF THE TNF- α -INDUCED RELEASE OF INTERLEUKIN 8 BY ENDOTHELIAL CELLS

[0141] It is well known to the person skilled in the art that the release of interleukin 8 (IL8) causes a massive accumulation of activated neutrophils and thus participates in the inflammatory process and/or tissue degradation process (see Baggiolini M. *et al.*; FEBS Letters, 307, 1, 97-101, 1992).

[0142] Human endothelial cells are grown at 37°C in multi-well plates under a water-saturated atmosphere constituted of a gaseous mixture of 95% air and 5% CO₂. Their culture medium is constituted by a medium M 199 pH = 7.4 containing 20% foetal calf serum, 2 mM L-glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin and 1% by volume of a medium supplement containing heparin and a growth factor for these cells.

[0143] When the cells are close to confluence, they are incubated for one hour in the presence or in the absence of one of the tested compounds of formula (I). This compound of the invention is incorporated at 5 µM in the culture medium containing 2% foetal calf serum, everything else being equal. After removal of the culture medium, the cells are incubated in the presence or in the absence (control) of TNF- α , at 1 ng/ml, in the same culture medium as before. In the case of cells pre-treated with a compound of the invention, the medium further contains the same compound at 5 µM. After three hours and thirty minutes of incubation, the interleukin 8 (IL-8) released into the culture medium is determined by ELISA.

[0144] The results obtained are given in the annexed Figure 1. These results show that the incubation of the endothelial cells in the presence of TNF- α leads to an increase in the production of IL-8 in the culture medium, and that the treatment of the cells by the compounds of formula (I) inhibits this effect by at least 70%.

[0145] These results demonstrate that such compounds can act as TNF- α antagonists in terms of interleukin 8 release by endothelial cells.

EXAMPLE 22: INHIBITION OF TNF- α -INDUCED P- AND E-SELECTIN EXPRESSIONS BY ENDOTHELIAL CELLS

[0146] It has been shown that the initial phase of inflammation is mediated by adhesion molecules of the E- and/or P-selectin type (see ALBELDA S.M. *et al.*, FASEB Journal, 8, 504-512, 1994).

[0147] Human endothelial cells are grown under the same conditions as those described in Example 21.

[0148] When the cells are close to confluence, they are incubated for one hour in the presence or in the absence of one of the tested compounds of formula (I). This compound of the invention is incorporated at 5 µM in the culture medium containing 2% foetal calf serum, everything else being equal. After the removal of the culture medium, the cells are incubated in the presence or in the absence (control) of TNF- α , at 1 ng/ml, in the same culture medium as before. In the case of cells pre-treated with a compound, the medium further contains the same compound at 10 µM. After three hours and thirty minutes of incubation, the cells are washed with PBS buffer and they are fixed with 2%

formaldehyde in the same buffer. The P- and E- selectin expressions on the cells are then measured by an ELISA determination by successively incubating the cells in the presence of a mouse monoclonal antibody, anti-P-selectin and anti-E-selectin respectively, and a rabbit anti-mouse anti-antibody labelled with alkaline phosphatase. The quantification is carried out upon the addition of para-nitrophenyl phosphate whose hydrolysis is followed at 405 nm.

[0149] The results obtained corresponding to the measurements of P- and E-selectin expressions are shown in the annexed Figures 2 and 3 respectively. These results show that the incubation of endothelial cells in the presence of TNF- α induces the expression of P- and E-selectin, which is inhibited by at least 75% and 90%, respectively, when the cells are treated with the tested compounds of formula (I).

[0150] The results show that such compounds are capable of inhibiting the TNF- α -induced expression of cell adhesion molecules such as P- and E-selectin.

[0151] The whole of these results show that the compounds of the invention having formula I:

1/ catalyse the reduction of hydroperoxides in the presence of glutathione;


2/ antagonize the action of TNF- α ;

3/ inhibit the expression of cell adhesion molecules.

[0152] TNF- α , as well as the expression of adhesion molecules such as P- and E-selectins, having been implicated in pathologies caused by an over-production of hydroperoxides, the molecules of the present invention of the general formula I therefore constitute valuable active principles and, after formulation, powerful drugs enabling the treatment of the corresponding pathologies.

Claims

1. Organoselenium compounds of general formula (I):



in which :

R = hydrogen ; -C(R₁R₂)-A-B ;

R₁ = lower alkyl, straight or branched, having from 1 to 8 carbon atoms ; optionally substituted aryl ; optionally substituted lower aralkyl ; wherein the term "aryl" means an aromatic group selected from phenyl and naphthyl ; wherein the term "substituted" concerning the terms "aryl", "aralkyl" means that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonyl, sulphonamide, sulphonyl(C₁-C₈)alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms ; and wherein "lower aralkyl" means phenyl(C₁-C₈) alkyl or naphthyl(C₁-C₈) alkyl and heteroar(C₁-C₈) alkyl ;


R₂ = lower alkyl, straight or branched, having from 1 to 8 carbon atoms ; optionally substituted aryl ; optionally substituted lower aralkyl ; wherein the term "aryl" means an aromatic group selected from phenyl and naphthyl ; wherein the term "substituted" concerning the terms "aryl", "aralkyl" means that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonyl, sulphonamide, sulphonyl(C₁-C₈)alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms ;

A = CO ; (CR₃R₄)_n

B represents NR₅R₆ ; N⁺R₅R₆R₇Y⁻ OR₅ ; SR₅ ;

Ar = an optionally substituted phenyl group or an optionally substituted radical of formula:


5



10

in which Z represents O ; S ; NR₅ ; when R = -C(R₁R₂)-A-B
or
Ar = a radical of formula

15



20


in which Z = O ; S ; NR₅ ; when R is hydrogen ;
wherein "substituted" concerning the terms "phenyl", "radical" (5-membered, including Z) means that the groups are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonyl, sulphonamide, sulpho(C₁-C₈)alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms,
X = Ar(R)-Se- ; -S-glutathione ; -S-N-acetylcysteine ; -S-cysteine ; -S-penicillamine ; -S-albumin ; -S-glucose ;

30

35

40

45



50

55

R₃ = hydrogen ; lower alkyl ; straight or branched, having from 1 to 8 carbon atoms ; optionally substituted aryl ; optionally substituted lower aralkyl ; wherein the term "aryl" means an aromatic group selected from phenyl and naphthyl ; wherein the term "substituted" concerning the terms "aryl", "aralkyl" means that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonyl, sulphonamide, sulpho(C₁-C₈)alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms, and wherein "lower aralkyl" means phenyl(C₁-C₈) alkyl or naphthyl(C₁-C₈) alkyl and heteroar (C₁-C₈) alkyl ;

R₄ = hydrogen ; lower alkyl ; straight or branched, having from 1 to 8 carbon atoms ; optionally substituted aryl ; optionally substituted lower aralkyl ; wherein the term "aryl" means an aromatic group selected from


phenyl and naphthyl; wherein the term "substituted" concerning the terms "aryl", "aralkyl" means that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonamide, sulpho(C₁-C₈)alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms, and wherein "lower aralkyl" means phenyl(C₁-C₈) alkyl or naphthyl(C₁-C₈) alkyl and heteroar(C₁-C₈) alkyl;

R₅ = hydrogen ; lower alkyl ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted heteroaryl ; optionally substituted lower heteroaralkyl ; CO(lower alkyl) ; CO(aryl) ; SO₂ (lower alkyl) ; SO₂(aryl) wherein "lower" in "lower alkyl" or "lower aralkyl", or "lower heteroaralkyl", or in "CO(lower alkyl)" or in "SO₂(lower alkyl)" means an alkyl group, straight or branched, having from 1 to 8 carbon atoms; the term "aryl" means an aromatic group selected from phenyl and naphthyl groups, the term "heteroaryl" means a mono- or bicyclic aromatic group, each cycle, or ring, comprising five or six atoms and said cycle, or ring, or both cycles, or rings, including in its carbon skeleton from one to three heretoatoms selected from nitrogen, oxygen and sulphur; and the term "substituted" concerning the terms "aryl", "aralkyl", "heteroaryl", "heteroaralkyl" means that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonamide, sulpho(C₁-C₈)alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms; the terms "lower aralkyl" and "lower heteroaralkyl" means phenyl(C₁-C₈) alkyl or naphthyl(C₁-C₈) alkyl and heteroar(C₁-C₈) alkyl respectively;

R₆ = hydrogen ; lower alkyl, straight or branched having from 1 to 8 carbon atoms; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted heteroaryl ; optionally substituted lower heteroaralkyl ; the term "aryl" means an aromatic group selected from phenyl and naphthyl groups, the term "heteroaryl" means a mono- or bicyclic aromatic group, each cycle, or ring, comprising five or six atoms and said cycle, or ring, or both cycles, or rings, including in its carbon skeleton from one to three heretoatoms selected from nitrogen, oxygen and sulphur; the term "substituted" concerning the terms "aryl, aralkyl, heteroaryl, heteroaralkyl" meaning that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonamide, sulpho(C₁-C₈) alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms, the terms "lower aralkyl" and "lower heteroaralkyl" meaning phenyl (C₁-C₈) alkyl or naphthyl (C₁-C₈) alkyl and heteroar (C₁-C₈) alkyl respectively ;

R₇ = hydrogen ; lower alkyl, straight or branched having from 1 to 8 carbon atoms ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted heteroaryl ; optionally substituted lower heteroaralkyl ; the term "aryl" means an aromatic group selected from phenyl and naphthyl groups, the term "heteroaryl" means a mono- or bicyclic aromatic group, each cycle, or ring, comprising five or six atoms and said cycle, or ring, or both cycles, or rings, including in its carbon skeleton from one to three heretoatoms selected from nitrogen, oxygen and sulphur; the term "substituted" concerning the terms "aryl, aralkyl, heteroaryl, heteroaralkyl" meaning that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonamide, sulpho(C₁-C₈) alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms, the terms "lower aralkyl" and "lower heteroaralkyl" meaning phenyl (C₁-C₈) alkyl or naphthyl (C₁-C₈) alkyl and heteroar (C₁-C₈) alkyl respectively ;

R₈ = hydrogen ; trifluoromethyl; lower alkyl ; straight or branched having from 1 to 8 carbon atoms ; optionally substituted aryl ; optionally substituted lower aralkyl ; optionally substituted heteroaryl ; optionally substituted lower heteroaralkyl ; the term "aryl" means an aromatic group selected from phenyl and naphthyl groups, the term "heteroaryl" means a mono- or bicyclic aromatic group, each cycle, or ring, comprising five or six atoms and said cycle, or ring, or both cycles, or rings, including in its carbon skeleton from one to three heretoatoms selected from nitrogen, oxygen and sulphur; the term "substituted" concerning the terms "aryl, aralkyl, heteroaryl, heteroaralkyl" meaning that the groups in question are substituted on the aromatic part with one or more identical or different groups selected from the groups : (C₁-C₈)alkyl, trifluoromethyl, (C₁-C₈)alkoxy, hydroxy, nitro, amino, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, sulphonyl, sulphonamide, sulpho(C₁-C₈) alkyl, carboxyl, carbalkoxyl, carbamide (it being possible for said (C₁-C₈)alkyl groups to be linear or branched) or substituted with one or more halogen atoms, the terms "lower aralkyl" and "lower heteroaralkyl" meaning phenyl (C₁-C₈) alkyl or naphthyl (C₁-C₈) alkyl and heteroar (C₁-C₈) alkyl respectively;



$n = 0$ or 1 ;

X^+ represents the cation of a pharmaceutically acceptable base ;

Y^- represents the anion of a pharmaceutically acceptable acid ;

and their salts of pharmaceutically acceptable acids or bases ;

with the provisos that:

when

$R = -C(R_1R_2)-(CR_3R_4)-B$ with $B = NR_5R_6$ or $N^+R_5R_6R_7Y^-$

and

$X = Ar(R)-Se-$ with $Ar =$ optionally substituted phenyl,

then

$-C(R_1R_2)$ is different from (CR_3R_4) ;

and

when

$Ar =$ phenyl and

$R = -C(R_1R_2)-C(O)-B$ with $B = NH_2$ or $NHCH_3$ or $NHCH_2C_6H_5$ or NHC_6H_5

and

$X = Ar(R)-Se-$,

then R_1 and R_2 cannot simultaneously represent a methyl group;

2. An organoselenium compound of general formula (I), according to claim 1, selected from the group consisting in:

- the di[2-[2'-(1'-amino-2'-methyl)propyl]phenyl]-diselenide,
- the di[2-[2'-(1'-amino-2'-methyl)propyl]phenyl]-diselenide dihydrochloride,
- the di[2-[2'-(1'-ammonium-2'-methyl)propyl]phenyl]-diselenide di-paratoluenesulphonate,
- the di[2-[2'-(1'-amino-2'-methyl)propyl]-4-methoxy]phenyl-diselenide,
- the di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenide,
- the di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenide dihydrochloride,
- the di[2-[2'-(1'-dimethylamino-2'-methyl)propyl]-phenyl]-diselenide,
- the di[2-[2'-(1'-trimethylammonium-2'-methyl)propyl]phenyl]-diselenide di-paratoluenesulphonate,
- the di [3-[2'-(1'-amino-2'-methyl) propyl]-2-thienyl] diselenide,
- the di [2-[2'-(1'-amino-2'-methyl) propyl]-3-thienyl]-diselenide,
- the S-(N-acetyl-L-cysteinyl)-[2-[2'-(1'-amino-2'-methyl)-propyl]-phenyl]-selenide,

- the S-glutathionyl-[2-[2'-(1'-amino-2'-methyl)-propyl]-phenyl]-selenide,
- the bis-4(5)-[2-phenyl-4(5)-seleno-1H-imidazole],
- the bis-4(5)-[[2-(4'-carbomethoxy)phenyl]-4(5)-seleno-1H-imidazole],
- the bis-4(5)-[2-[(4'-carboxyl)phenyl]-4(5)-seleno-1H-imidazole],
- the bis-4(5)-[2-[4'-carbo(2''-(4'''-methyl-piperazin-1''-yl)ethoxyphenyl)]-4(5)-seleno-1H-imidazole],
- the bis-5-[2-[(4'-trifluoromethyl)phenyl]-5-seleno-1-methyl-imidazole],
- the bis-5-[2-[(4'-carbomethoxy)phenyl]-5-seleno-1-methyl-imidazole],
- the bis-5-[2-[(4'-carboxyl)-phenyl]-5-seleno-1-methyl-imidazole].

3. Use of organoselenium compounds, of the general formula (I),

according to claim 1 or 2, as antioxidants ;

intended in particular to be added to preserving media of grafts for transplantation of organs of human or animal origin such as the heart, the liver, the kidney and the lungs;

and

in particular for the manufacture of pharmaceutical compositions with antioxidant activity, intended especially :

- * for treatments of any physiopathological condition in which an over-production of cytotoxic hydroperoxides contributes to the functional impairments of cells or tissues ; and more particularly including :
- * the treatment of inflammatory and/or ischaemic cardio- and cerebro-vascular pathologies, such as the preventive and/or curative treatment of arterial restenoses following an angioplasty, the preventive and/or curative treatment of arterial stenoses following artery allografts, the treatment of intermittent claudication in patients affected with obstructive ischaemia of the lower members, the treatment of cerebro-vascular accidents of ischaemic origin ;
- * the treatment of inflammatory and/or ischaemic digestive pathologies, such as the treatment of acute inflammations of the bowel (Crohn's disease, hemorrhagic rectocolitis) ;
- * the treatment of inflammatory and/or ischaemic respiratory pathologies, such as the treatment of adult respiratory distress syndrome (ARDS) and infant respiratory distress syndrome (IRDS) ;
- * the treatment of inflammatory and/or ischaemic ophthalmic pathologies, such as the treatment of glaucoma ;
- * the treatment of cataracts ;
- * the treatment of acute ophthalmic allergies ;
- * the treatment of impairments of the retina which are associated with a macular degeneration;
- * the treatment of viral infections causing an immuno-deficiency, such as the treatment of AIDS ; and
- * the treatment of post-radiotherapy fibroses.

4. Use of organo selenium compounds, of the general formula (I), according to claim 1 or 2, for the manufacture of pharmaceutical compositions intended for the treatment:

- * of inflammatory diseases of the bowel such as Crohn's disease or hemorrhagic rectocolitis ;
- * of adult respiratory distress syndrome and infant respiratory distress syndrome ;
- * of cataracts ;
- * of AIDS ;
- * of post-radiotherapy fibroses.

5. A pharmaceutical composition, notably having an antioxidant activity, characterized in that it comprises, as active ingredient, at least one organoselenium compound, of the general formula (I), according to claim 1 or 2, or one of its pharmaceutically acceptable salts of an acid or a base, optionally incorporated in a pharmaceutically acceptable excipient, carrier or vehicle.

6. The pharmaceutical composition according to claim 5, characterised in that said organoselenium compound of the general formula (I) above, optionally as salt, is present in an amount between 0.1 and 5 % by weight, advantageously between 0.1 and 1 % by weight, based on the total weight of the composition.

7. The pharmaceutical composition according to one of claims 5 or 6, characterized in that it is in the form of a unit dose comprising from 1 to 500 mg of at least one organoselenium compound of the general formula (I).

8. The pharmaceutical composition according to any one of claims 5 to 7, characterized in that it is a pharmaceutical

composition for the treatment of any physiopathological condition in which an over-production of cytotoxic hydroperoxides contributes to the functional impairments of cells or tissues ; this over-production of hydroperoxide can be due :

- 5 - to the activation of the intra-cellular metabolic pathways such as those of the flavine or cytochrome P-450 oxygenases, those of the lipoxygenases, and those of the monoamine oxidases ;
- to the activation of enzymes contained in endothelial cells (xanthine oxidase, 15-lipoxygenase) or in blood platelets (cyclooxygenase and 12-lipoxygenase) ;
- 10 - to the activation, by cytokines such as TNF- α , of inflammatory and/or immune cells such as neutrophils, macrophages or lymphocytes ;
- to an intoxication by a free-radical generating xenobiotic ;
- to a voluntary irradiation, such as practised during a radiotherapy, or an accidental irradiation.

9. The pharmaceutical composition according to any one of claims 5 to 6, characterized in that it is a pharmaceutical composition for the treatment of any physiopathological condition in which an over-production of cytotoxic hydroperoxides contributes to the functional impairments of cells or tissues ; and more particularly :

- * for the treatment of inflammatory and/or ischaemic cardio- and cerebro-vascular pathologies, such as the preventive and/or curative treatment of arterial restenoses following an angioplasty, the preventive and/or curative treatment of arterial stenoses following artery allografts, the treatment of intermittent claudication in patients affected with obstructive ischaemia of the lower members, the treatment of cerebro-vascular accidents of ischaemic origin ;
- 20 * for the treatment of inflammatory and/or ischaemic digestive pathologies, such as the treatment of acute inflammations of the bowel (Crohn's disease, hemorrhagic rectocolitis) ;
- 25 * for the treatment of inflammatory and/or ischaemic respiratory pathologies, such as the treatment of adult respiratory distress syndrome (ARDS) and infant respiratory distress syndrome (IRDS) ;
- * for the treatment of inflammatory and/or ischaemic ophthalmic pathologies, such as the treatment of glaucoma ;
- * for the treatment of cataracts ;
- 30 * for the treatment of acute ophthalmic allergies ;
- * for the treatment of impairments of the retina which are associated with a macular degeneration ;
- * for the treatment of viral infections causing an immuno-deficiency, such as the treatment of AIDS ; and
- * for the treatment of post-radiotherapy fibroses.

35 10. Method of preparation of organoselenium compounds according to claim 1 or 2, of general formula (I), in which R = -C(R₁R₂)-A-B (R \neq H) ; characterised in that it comprises the following essential steps :

a/ preparing or using an orthohalo(hetero)arylacetonitrile derivative, optionally mono-or gem- disubstituted in the benzylic position; then, according to the series considered :

- 40 - for the preparation of said compounds of formula (I) in which A=(CR₃R₄)_n with n = 0 (A does not exist) :

b1/ hydrolyzing said nitrile derivative into an amide derivative,

45 c1-1/ transforming this into an amine derivative by a transposition reaction according to conventional methods,

d1-1/ allowing said amine derivative to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylisoselenazoline derivative,

50 e1-1/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, according to conventional procedures, said (hetero)arylisoselenazoline derivative ;

- for the preparation of said compounds of formula (I) in which A = CO:

b1/ hydrolyzing said nitrile derivative into an amide derivative,

55 c1-2/ allowing said amide derivative to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylisoselenazone derivative,

d1-2/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, said (hetero)arylisose-

lenazone derivative;

- for the preparation of said compounds of formula (I) in which $A = CH_2$:

5 b2/ reducing said nitrile derivative into an amine derivative, with the aid of borane for example in an etheral solvent such as tetrahydrofuran for example,
 c2/ allowing said amine derivative to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylselenazine derivative,
 10 d2/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, according to conventional procedures, said (hetero)arylselenazine derivative;


- for the preparation of said compounds of formula (I) in which $A = (CR_3R_4) (\neq CH_2)$:

15 b3/ carrying out a mono- or a bis-C-alkylation of said nitrile derivative according to conventional methods, with the aid of an organolithium derivative for example in an etheral solvent such as tetrahydrofuran;
 c3/ allowing the amine derivative obtained to react with a nucleophilic selenium derivative, optionally generated *in situ*, in the presence of a copper Cu(I) salt, in a polar organic solvent, to lead to the corresponding (hetero)arylselenazine derivative;
 20 d3/ optionally, N-alkylating or N-aryllating or N-acylating or N-sulphonylating, according to conventional procedures, said corresponding (hetero)arylselenazine derivative;

25 b/ reducing, in a polar solvent such as methanol, the cyclic compound obtained, according to any one of the above synthetic routes, with the aid of a metal hydride, such as sodium borohydride, intervening in an amount corresponding to a half-reducing equivalent; for the preparation of said compounds of formula (I) in which $X = Ar(R)-Se-$;

30 or
 allowing said cyclic compound obtained according to any one of the above synthetic routes to react at ambient temperature, with the thiol compound corresponding to the values of $X \neq Ar(R)-Se-$, for the preparation of said compounds of formula (I) in which $X \neq Ar(R)-Se-$.


35 11. Method of preparation of organoselenium compounds according to claim 1 or 2, of general formula (I) in which $R = H$; i.e. of general formula (II) below:



50 characterised in that it comprises the following essential steps:

- a/ preparing or using a derivative of formula (III):

55



10 N-alkylated, N-arylated, N-acylated or N-sulphonylated when $Z = NR_5$; then, according to the series considered :

- for the preparation of said diselenide compounds of formula (II) in which R_8 = hydrogen :

15 b1/ allowing said derivative of formula (III) to react with an electrophilic selenium derivative, in a non-polar solvent;

c1/ if necessary, de-acylating or de-sulphonylating, according to usual procedures, the compound obtained ;

- for the preparation of said diselenide compounds of formula (II) in which R_8 = lower alkyl, optionally substituted lower aralkyl, optionally substituted lower heteroaralkyl :

25 b2/ alkylating said derivative of formula (III) in position 2 in treating it firstly with an organolithium base such as lithium diisopropylamide, then, with a halide, notably a lower alkyl, optionally substituted lower aralkyl or optionally substituted lower heteroaralkyl iodide ;

c2/ allowing said alkyl derivative to react with an electrophilic selenium derivative, in a non-polar solvent;

d2/ if necessary, de-acylating or de-sulphonylating, according to usual procedures, the compound obtained ;

- for the preparation of said diselenide compounds of formula (II) in which R_8 = optionally substituted aryl, optionally substituted heteroaryl:

35 b3/ treating said derivative of formula (III), in a non-polar organic solvent, in the presence of a strong base, with a trialkyltin halide or a zinc halide in order to obtain the corresponding stannyl or zinc derivative then treating said stannyl or zinc derivative with a haloaromatic derivative, in the presence of palladium, in a non-polar organic solvent;


c3/ allowing the compound obtained to react with an electrophilic selenium derivative, in a non-polar solvent;

40 d3/ if necessary, de-acylating or de-sulphonylating, according to usual procedures, the compound obtained ;

45 b/ if necessary, allowing said diselenide compound obtained according to any one of the above synthetic routes to react, in a polar solvent, with an adequate mercaptan ; in order to obtain the corresponding selenosulphide compounds.

Patentansprüche

- 50 1. Organoselenverbindungen der allgemeinen Formel (I) :



worin:

R = Wasserstoff; $-C(R_1R_2)-A-B$;


R_1 = nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Alkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl und Naphthyl; wobei der Begriff "substituiert" betreffend die Begriffe "Aryl", "Alkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C_1-C_8) Alkyl, Trifluormethyl, (C_1-C_8) Alkoxy, Hydroxy, Nitro, Amino, (C_1-C_8) Alkylamino, Di (C_1-C_8) alkylamino, Sulfoxyl, Sulfonyl, Sulfonamid, Sulfo (C_1-C_8) alkyl, Carboxyl, Carbalkoxyl, Carbamid (wobei besagte (C_1-C_8) Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen, und worin "nied.-Alkyl" Phenyl (C_1-C_8) alkyl oder Naphthyl (C_1-C_8) alkyl und Heteroar (C_1-C_8) alkyl bedeutet;

R_2 = nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Alkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl und Naphthyl; wobei der Begriff "substituiert" betreffend die Begriffe "Aryl", "Alkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C_1-C_8) Alkyl, Trifluormethyl, (C_1-C_8) Alkoxy, Hydroxy, Nitro, Amino, (C_1-C_8) Alkylamino, Di (C_1-C_8) alkylamino, Sulfoxyl, Sulfonyl, Sulfonamid, Sulfo (C_1-C_8) alkyl, Carboxyl, Carbalkoxyl, Carbamid (wobei besagte (C_1-C_8) Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen;


A = CO; $(CR_3R_4)_n$;

B für NR_5R_6 ; $N^+R_5R_6R_7Y^-$; OR_5 ; SR_5 steht;

Ar = eine gegebenenfalls substituierte Phenylgruppe oder einen gegebenenfalls substituierten Rest der Formel:




darstellt, worin Z für O; S; NR_5 steht; wenn R = $-C(R_1R_2)-A-B$ oder Ar = ein Rest der Formel



worin Z = C; S; NR_5 ; wenn R Wasserstoff darstellt;

worin "substituiert" betreffend die Begriffe "Phenyl", "Rest" (5-gliedrig einschliesslich Z) bedeutet, dass die Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C_1-C_8) Alkyl, Trifluormethyl, (C_1-C_8) Alkoxy, Hydroxy, Nitro, Amino, (C_1-C_8) Alkylamino, Di (C_1-C_8) alkylamino, Sulfoxyl, Sulfonyl, Sulfonamid, Sulfo (C_1-C_8) alkyl, Carboxyl, Carbalkoxyl, Carbamid (wobei besagte (C_1-C_8) Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen,

X = Ar(R)-Se; -S-Glutathion; -S-N-Acetylcystein; -S-Cystein; -S-Penicillamin; -S-Albumin; -S-Glucose;



R₃ = Wasserstoff; nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Aralkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl und Naphthyl; wobei der Begriff "substituiert" betreffend die Begriffe "Aryl", "Aralkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C₁-C₈)Alkyl, Trifluormethyl, (C₁-C₈)Alkoxy, Hydroxy, Nitro, Amino, (C₁-C₈)Alkylamino, Di(C₁-C₈)alkylamino, Sulfoxyl, Sulfonyl, Sulfonamid, Sulfo(C₁-C₈)alkyl, Carboxyl, Carbalkoxyl, Carbamid (wobei besagte (C₁-C₈)Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen, und worin "nied.-Aralkyl" Phenyl(C₁-C₈)alkyl oder Naphthyl(C₁-C₈)alkyl und Heteroar(C₁-C₈)alkyl bedeutet;

R₄ = Wasserstoff; nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Aralkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl und Naphthyl; wobei der Begriff "substituiert" betreffend die Begriffe "Aryl", "Aralkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C₁-C₈)Alkyl, Trifluormethyl, (C₁-C₈)Alkoxy, Hydroxy, Nitro, Amino, (C₁-C₈)Alkylamino, Di(C₁-C₈)alkylamino, Sulfoxyl, Sulfonyl, Sulfonamid, Sulfo(C₁-C₈)alkyl, Carboxyl, Carbalkoxyl, Carbamid (wobei besagte (C₁-C₈)Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen, und worin "nied.-Aralkyl" Phenyl(C₁-C₈)alkyl oder Naphthyl(C₁-C₈)alkyl und Heteroar(C₁-C₈)alkyl bedeutet;


R₅ = Wasserstoff; nied.-Alkyl; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Aralkyl; gegebenenfalls substituiertes Heteroaryl; gegebenenfalls substituiertes nied.-Heteroaralkyl; CO(nied.-Alkyl); CO(Aryl); SO₂(nied.-Alkyl); SO₂(Aryl); wobei "nied." in "nied.-Alkyl" oder "nied.-Aralkyl" oder "nied.-Heteroaralkyl" oder in "CO(nied.-Alkyl)" oder in "SO₂(nied.-Alkyl)" eine Alkyl-Gruppe, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen bedeutet; der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl- und Naphthyl-Gruppen; der Begriff "Heteroaryl" eine mono- oder bicyclische aromatische Gruppe bedeutet, wobei jeder Kreis oder Ring fünf oder sechs Atome umfasst und besagter Kreis oder Ring oder beide Kreise oder Ringe, in ihrem Kohlenstoffgerüst ein bis drei Heteroatome, ausgewählt aus Stickstoff, Sauerstoff und Schwefel einschließen; und der Begriff "substituiert" betreffend die Begriffe "Aryl", "Aralkyl", "Heteroaryl", "Heteroaralkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C₁-C₈)Alkyl, Trifluormethyl, (C₁-C₈)Alkoxy, Hydroxy, Nitro, Amino, (C₁-C₈)Alkylamino, Di(C₁-C₈)alkylamino, Sulfoxyl, Sulfonyl, Sulfonamid, Sulfo(C₁-C₈)alkyl, Carboxyl, Carbalkoxyl, Carbamid (wobei besagte (C₁-C₈)Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen; die Begriffe "nied.-Aralkyl" und "nied.-Heteroaralkyl" Phenyl(C₁-C₈)alkyl oder Naphthyl(C₁-C₈)alkyl bzw. Heteroar(C₁-C₈)alkyl bedeuten;

R₆ = Wasserstoff; nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Aralkyl; gegebenenfalls substituiertes Heteroaryl; gegebenenfalls substituiertes nied.-Heteroaralkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl- und Naphthyl-Gruppen, der Begriff "Heteroaryl" eine mono- oder bicyclische aromatische Gruppe bedeutet, wobei jeder Kreis oder Ring fünf oder sechs Atome umfasst und besagter Kreis oder Ring oder beide Kreise oder Ringe, in ihrem Kohlenstoffgerüst ein bis drei Heteroatome, ausgewählt aus Stickstoff,

Sauerstoff und Schwefel einschliessen; der Begriff "substituiert" betreffend die Begriffe "Aryl", "Aralkyl", "Heteroaryl", "Heteroaralkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C₁-C₈) Alkyl, Trifluormethyl, (C₁-C₈) Alkoxy, Hydroxy, Nitro, Amino, (C₁-C₈) Alkylamino, Di(C₁-C₈) alkylamino, Sulfoxy, Sulfonoyl, Sulfonamid, Sulfo(C₁-C₈) alkyl, Carboxyl, Carbalkoxy, Carbamid (wobei besagte (C₁-C₈) Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen; die Begriffe "nied.-Aralkyl" und "nied.-Heteroaralkyl" Phenyl(C₁-C₈) alkyl oder Naphthyl(C₁-C₈) alkyl bzw. Heteroar(C₁-C₈) alkyl bedeuten;

R₇ = Wasserstoff; nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Aralkyl; gegebenenfalls substituiertes Heteroaryl; gegebenenfalls substituiertes nied.-Heteroaralkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl- und Naphthyl-Gruppen, der Begriff "Heteroaryl" eine mono- oder bicyclische aromatische Gruppe bedeutet, wobei jeder Kreis oder Ring fünf oder sechs Atome umfasst und besagter Kreis oder Ring oder beide Kreise oder Ringe, in ihrem Kohlenstoffgerüst ein bis drei Heteroatome, ausgewählt aus Stickstoff, Sauerstoff und Schwefel einschliessen; der Begriff "substituiert" betreffend die Begriffe "Aryl", "Aralkyl", "Heteroaryl", "Heteroaralkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C₁-C₈) Alkyl, Trifluormethyl, (C₁-C₈) Alkoxy, Hydroxy, Nitro, Amino, (C₁-C₈) Alkylamino, Di(C₁-C₈) alkylamino, Sulfoxy, Sulfonoyl, Sulfonamid, Sulfo(C₁-C₈) alkyl, Carboxyl, Carbalkoxy, Carbamid (wobei besagte (C₁-C₈) Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehr Halogenatomen; die Begriffe "nied.-Aralkyl" und "nied.-Heteroaralkyl" Phenyl(C₁-C₈) alkyl oder Naphthyl(C₁-C₈) alkyl bzw. Heteroar(C₁-C₈) alkyl bedeuten;

R₈ = Wasserstoff; Trifluormethyl; nied.-Alkyl, gerade oder verzweigt, mit 1 bis 8 Kohlenstoffatomen; gegebenenfalls substituiertes Aryl; gegebenenfalls substituiertes nied.-Aralkyl; gegebenenfalls substituiertes Heteroaryl; gegebenenfalls substituiertes nied.-Heteroaralkyl; wobei der Begriff "Aryl" eine aromatische Gruppe bedeutet, ausgewählt aus Phenyl- und Naphthyl-Gruppen, der Begriff "Heteroaryl" eine mono- oder bicyclische aromatische Gruppe bedeutet, wobei jeder Kreis oder Ring fünf oder sechs Atome umfasst und besagter Kreis oder Ring oder beide Kreise oder Ringe, in ihrem Kohlenstoffgerüst ein bis drei Heteroatome, ausgewählt aus Stickstoff, Sauerstoff und Schwefel einschliessen; der Begriff "substituiert" betreffend die Begriffe "Aryl", "Aralkyl", "Heteroaryl", "Heteroaralkyl" bedeutet, dass die fraglichen Gruppen an dem aromatischen Teil mit einer oder mehreren identischen oder unterschiedlichen Gruppen substituiert sind, ausgewählt aus den Gruppen: (C₁-C₈) Alkyl, Trifluormethyl, (C₁-C₈) Alkoxy, Hydroxy, Nitro, Amino, (C₁-C₈) Alkylamino, Di(C₁-C₈) alkylamino, Sulfoxy, Sulfonoyl, Sulfonamide, Sulfo(C₁-C₈) alkyl, Carboxyl, Carbalkoxy, Carbamid (wobei besagte (C₁-C₈) Alkylgruppen gerade oder verzweigt sein können) oder substituiert mit einem oder mehreren Halogenatomen; die Begriffe "nied.-Aralkyl" und "nied.-Heteroaralkyl" Phenyl(C₁-C₈) alkyl oder Naphthyl(C₁-C₈) alkyl bzw. Heteroar(C₁-C₈) alkyl bedeuten;



n = 0 oder 1;

X⁺ das Kation einer pharmazeutisch annehmbaren Base darstellt;

Y⁻ das Anion einer pharmazeutisch annehmbaren Säure darstellt; und ihre Salze von pharmazeutisch an-

nehmbaren Säuren oder Basen; unter den Bedingungen, dass:

wenn

- 5 $R = -C(R_1R_2)-(CR_3R_4)-B$ mit $B = NR_5R_6$ oder $N^+R_5R_6R_7Y^-$ und
 $X = Ar(R)-Se$ mit $Ar =$ gegebenenfalls substituiertes Phenyl, dann
 $-C(R_1R_2)$ unterschiedlich ist von (CR_3R_4) ; und

wenn

- 10 $Ar =$ Phenyl und
 $R = -C(R_1R_2)-C(O)-B$ mit $B = NH_2$ oder $NHCH_3$ oder $NHCH_2C_6H_5$ oder NHC_6H_5 und
 $X = Ar(R)-Se$, dann können R_1 und R_2 nicht gleichzeitig eine Methyl-Gruppe darstellen.

- 15 2. Organoselenverbindung der allgemeinen Formel (I) gemäss Anspruch 1, ausgewählt aus der Gruppe bestehend aus:

- dem Di[2-[2'-(1'-amino-2'-methyl)propyl]phenyl]-diselenid,
- dem Di[2-[2'-(1'-amino-2'-methyl)propyl]phenyl]-diselenid-dihydrochlorid,
- 20 - dem Di[2-[2'-(1'-ammonium-2'-methyl)propyl]phenyl]-diselenid-diparatoluolsulfonat,
- dem Di[2-[2'-(1'-amino-2'-methyl)propyl]-4-methoxyphenyl]-diselenid,
- dem Di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenid,
- dem Di[2-[2'-(1'-methylamino-2'-methyl)propyl]-phenyl]-diselenid-dihydrochlorid,
- dem Di[2-[2'-(1'-dimethylamino-2'-methyl)propyl]-phenyl]-diselenid,
- 25 - dem Di[2-[2'-(1'-trimethylammonium-2'-methyl)propyl]phenyl]-diselenid-di-paratoluolsulfonat,
- dem Di[3-[2'-(1'-amino-2'-methyl)propyl]-2-thienyl]-diselenid,
- dem Di[2-[2'-(1'-amino-2'-methyl)propyl]-3-thienyl]-diselenid,
- dem S-(N-acetyl-L-cysteinyl)-[2-[2'-(1'-amino-2'-methyl)propyl]-phenyl]-selenid,
- dem S-Glutathionyl-[2-[2'-(1'-amino-2'-methyl)propyl]-phenyl]-selenid,
- 30 - dem Bis-4(5)-[2-phenyl-4(5)-seleno-1H-imidazol],
- dem Bis-4(5)-[[2-(4'-carbomethoxyphenyl)-4(5)-seleno-1H-imidazol],
- dem Bis-4(5)-[2-[(4'-carboxyl)phenyl]-4(5)-seleno-1H-imidazol],
- dem Bis-4(5)-[2-[4'-(2"-methyl-piperazin-1"-yl)ethoxyphenyl]-4(5)-seleno-1H-imidazol],
- dem Bis-5-[2-[(4'-trifluormethyl)phenyl]-5-seleno-1-methylimidazol],
- 35 - dem Bis-5-[2-[(4'-carbomethoxyphenyl)-5-seleno-1-methylimidazol],
- dem Bis-5-[2-[(4'-carboxyl)-phenyl]-5-seleno-1-methylimidazol].

3. Verwendung von Organoselenverbindungen der allgemeinen Formel (I) gemäss Anspruch 1 oder 2 als Antioxidantien; beabsichtigt insbesondere um zu Schutzmitteln von Transplantaten für Organtransplantationen menschlichen oder tierischen Ursprungs wie dem Herzen, der Leber, den Nieren und den Lungen zugegeben zu werden; und insbesondere für die Herstellung von pharmazeutischen Zusammensetzungen mit Antioxidanz-Wirksamkeit, beabsichtigt besonders:

- * für Behandlungen jedes physiopathologischen Zustands, in welchem eine Überproduktion von cytotoxischen Hydroperoxiden zu den funktionellen Beeinträchtigungen von Zellen oder Geweben beiträgt; und insbesondere einschliessend:
- * die Behandlung von entzündlichen und/oder ischämischen kardio- und cerebrovaskulären Pathologien, wie die präventive und/oder kurative Behandlung arterieller Restenosen infolge einer Angioplastie, die präventive und/oder kurative Behandlung von arteriellen Stenosen infolge von Arterie-Allografts, die Behandlung von Claudicatio intermittens bei Patienten, welche von obstruktiver Ischämie der unteren Glieder betroffen sind, die Behandlung von cerebro-vaskulären Unfällen ischämischen Ursprungs;
- * die Behandlung von entzündlichen und/oder ischämischen digestiven Pathologien, wie die Behandlung akuter Entzündungen des Darms (Crohn-Krankheit, hämorrhagischer Rektokolitis);
- * die Behandlung von entzündlichen und/oder ischämischen Respirationspathologien, wie die Behandlung von "adult respiratory distress syndrome" (ARDS) und "infant respiratory distress syndrome" (IRDS);
- 55 * die Behandlung von entzündlichen und/oder ischämischen ophthalmologischen Pathologien, wie die Behandlung von Glaukom;
- * die Behandlung von Katarakten;

- * die Behandlung von akuten ophthalmologischen Allergien;
 - * die Behandlung von Beeinträchtigungen der Retina welche mit einer Makuladegeneration in Verbindung stehen;
- 5 die Behandlung von Virusinfektionen, welche eine Immunschwäche bewirken, wie die Behandlung von AIDS;
und
 die Behandlung von Post-Radiotherapie-Fibrosen.
- 10 4. Verwendung von Organoselenverbindungen der allgemeinen Formel (I) gemäss Anspruch 1 oder 2 für die Herstellung von pharmazeutischen Zusammensetzungen, beabsichtigt für die Behandlung:
- * von entzündlichen Erkrankungen des Darms wie Crohn-Krankheit oder hämorrhagischer Rektokolitis;
 - * von "adult respiratory distress syndrome" und "infant respiratory distress syndrome";
 - * von Katarakten;
 - 15 * von AIDS;
 - * von Post-Radiotherapie-Fibrosen.
- 20 5. Pharmazeutische Zusammensetzung, vorwiegend mit einer Antioxidanz-Wirksamkeit, dadurch gekennzeichnet, dass sie als Wirkstoff mindestens eine Organoselenverbindung der allgemeinen Formel (I) gemäss Anspruch 1 oder 2 umfasst, oder eine ihrer pharmazeutisch annehmbaren Salze einer Säure oder Base, gegebenenfalls eingeschlossen in einen pharmazeutisch annehmbaren Arzneimittelträger, Träger oder Vehikel.
- 25 6. Pharmazeutische Zusammensetzung gemäss Anspruch 5, dadurch gekennzeichnet, dass besagte Organoselenverbindung der allgemeinen Formel (I) oben, gegebenenfalls als Salz, in einer Menge zwischen 0,1 und 5 Gew.-%, vorteilhafterweise zwischen 0,1 und 1 Gew.-%, basierend auf dem Gesamtgewicht der Zusammensetzung vorhanden ist.
- 30 7. Pharmazeutische Zusammensetzung gemäss einem der Ansprüche 5 oder 6, dadurch gekennzeichnet, dass sie sich in der Form einer Einheitsdosis befindet, welche 1 bis 500 mg von mindestens einer Organoselenverbindung der allgemeinen Formel (I) umfasst.
- 35 8. Pharmazeutische Zusammensetzung gemäss einem der Ansprüche 5 bis 7, dadurch gekennzeichnet, dass sie eine pharmazeutische Zusammensetzung für die Behandlung von jedem physiopathologischen Zustand ist, in welchem eine Überproduktion von cytotoxischen Hydroperoxiden zu funktionellen Beeinträchtigungen von Zellen und Geweben beiträgt; diese Überproduktion von Hydroperoxid kann
- der Aktivierung der intrazellulären, metabolischen Wege, wie jene der Flavin- oder Cytochrom-P-450-Oxygenasen, jene der Lipoxygenasen und jene der Monoaminoxidasen;
 - der Aktivierung von Enzymen, welche in endothelialen Zellen (Xanthinoxidase, 15-Lipoxygenase) oder in Blutplättchen (Cyclooxygenase und 12-Lipoxygenase) enthalten sind
 - 40 - der Aktivierung durch Cytokine, wie TNF- α , von Entzündungs- und/oder Immunzellen wie Neutrophilen, Makrophagen oder Lymphozyten;
 - einer Intoxikation durch ein freie Radikale erzeugendes Xenobiotikum;
 - einer freiwilligen Bestrahlung, wie sie während einer Radiotherapie praktiziert wird, oder einer versehentlichen
 - 45 Bestrahlung
- zuzuschreiben sein.
- 50 9. Pharmazeutische Zusammensetzung gemäss einem der Ansprüche 5 bis 6, dadurch gekennzeichnet, dass es eine pharmazeutische Zusammensetzung für die Behandlung eines jeden physiopathologischen Zustands ist, in welchem eine Überproduktion von cytotoxischen Hydroperoxiden zu der funktionellen Beeinträchtigung von Zellen oder Geweben beiträgt; und insbesondere:
- 55 *
- für die Behandlung von entzündlichen und/oder ischämischen kardio- und cerebrovaskulären Pathologien, wie die präventive und/oder kurative Behandlung arterieller Restenosen infolge einer Angioplastie, die präventive und/oder kurative Behandlung von arteriellen Stenosen infolge von Arterie-Allografts, die Behandlung von Claudicatio intermittens bei Patienten, welche von obstruktiver Ischämie der unteren Glieder betroffen sind, die Behandlung von cerebro-vaskulären Unfällen ischämischen Ursprungs;

- * die Behandlung von entzündlichen und/oder ischämischen digestiven Pathologien, wie die Behandlung akuter Entzündungen des Darms (Crohn-Krankheit, hämorrhagischer Rektokolitis);
- * für die Behandlung von entzündlichen und/oder ischämischen Respirationspathologien, wie die Behandlung von "adult respiratory distress syndrome" (ARDS) und "infant respiratory distress syndrome" (IRDS);
- 5 * für die Behandlung von entzündlichen und/oder ischämischen ophthalmologischen Pathologien, wie die Behandlung von Glaukom;
- * für die Behandlung von Katarakten;
- * für die Behandlung von akuten ophthalmologischen Allergien;
- 10 * für die Behandlung von Beeinträchtigungen der Retina, welche mit einer Makuladegeneration in Verbindung stehen;

für die Behandlung von Virusinfektionen, welche eine Immunschwäche bewirken, wie die Behandlung von AIDS; und

für die Behandlung von Post-Radiotherapie-Fibrosen.

10. Verfahren zur Herstellung von Organoselenverbindungen gemäss Anspruch 1 oder 2 der allgemeinen Formel (I), worin $R = -C(R_1R_2)-A-B$ ($R \neq H$); dadurch gekennzeichnet, dass es die folgenden wesentlichen Schritte umfasst:

a/ Herstellen oder Verwenden eines Orthohalogen(hetero)arylacetonitrilderivats, gegebenenfalls mono- oder gem-disubstituiert in der Benzyl-Position; dann, gemäss der in Erwägung gezogenen Serie:

- für die Herstellung von besagten Verbindungen der Formel (I), worin $A = (CR_3R_4)_n$ mit $n = 0$ (A existiert nicht):

b1/ Hydrolisieren von besagtem Nitrilderivat zu einem Amidderivat,
 c1-1/ Umwandeln von diesem in ein Aminderivat durch eine Umsetzungsreaktion gemäss herkömmlicher Verfahren,
 d1-1/ Erlauben, dass sich das besagte Aminderivat mit einem nucleophilen Selenderivat umsetzt, gegebenenfalls erzeugt *in situ*, in Anwesenheit von einem Kupfer-Cu(I)-Salz in einem polaren organischen Lösungsmittel, um zu dem entsprechenden (Hetero)arylisoselenazolinderivat zu führen,
 e1-1/ gegebenenfalls N-Alkylieren oder N-Arylieren oder N-Acylieren oder N-Sulfonylieren des besagten (Hetero)arylisoselenazolinderivats gemäss herkömmlicher Verfahren,

- für die Herstellung von besagten Verbindungen der Formel (I), worin $A = CO$:

b1/ Hydrolisieren von besagtem Nitrilderivat zu einem Amidderivat,
 c1-2/ Erlauben, dass sich das besagte Amidderivat mit einem nucleophilen Selenderivat umsetzt, gegebenenfalls erzeugt *in situ*, in Anwesenheit von einem Kupfer-Cu(I)-Salz in einem polaren organischen Lösungsmittel, um zu dem entsprechenden (Hetero)arylisoselenazonderivat zu führen,
 d1-2/ gegebenenfalls N-Alkylieren oder N-Arylieren oder N-Acylieren oder N-Sulfonylieren des besagten (Hetero)arylisoselenazonderivats;

- für die Herstellung besagter Verbindungen der Formel (I), worin $A = CH_2$:

b2/ Reduzieren des besagten Nitrilderivats zu einem Aminderivat mit Hilfe von Boran beispielsweise in einem etherischem Lösungsmittel wie Tetrahydrofuran zum Beispiel,
 c2/ Erlauben, dass sich das besagte Aminderivat mit einem nucleophilen Selenderivat umsetzt, gegebenenfalls erzeugt *in situ*, in Anwesenheit von einem Kupfer-Cu(I)-Salz in einem polaren organischen Lösungsmittel, um zu dem entsprechenden (Hetero)arylisoselenazinderivat zu führen,
 d2/ gegebenenfalls N-Alkylieren oder N-Arylieren oder N-Acylieren oder N-Sulfonylieren des besagten (Hetero)arylisoselenazinderivats gemäss herkömmlicher Verfahren

- für die Herstellung besagter Verbindungen der Formel (I), worin $A = (CR_3R_4) (\neq CH_2)$:


b3/ Durchführen einer Mono- oder einer Bis-C-Alkylierung des besagten Nitrilderivats gemäss herkömmlicher Verfahren, mit Hilfe eines Organolithiumderivats beispielsweise in einem etherischen Lösungsmittel wie Tetrahydrofuran;
 c3/ Erlauben, dass sich das erhaltene Aminderivat mit einem nucleophilen Selenderivat umsetzt, ge-

gegebenenfalls erzeugt *in situ*, in Anwesenheit von einem Kupfer-Cu(I)-Salz in einem polaren organischen Lösungsmittel, um zu dem entsprechenden (Hetero)aryliselenazinderivat zu führen, d3/ gegebenenfalls N-Alkylieren oder N-Arylieren oder N-Acylieren oder N-Sulfonylieren des besagten entsprechenden (Hetero)aryliselenazinderivats gemäss herkömmlicher Verfahren;

b/ Reduzieren der gemäss einem der obigen synthetischen Wege erhaltenen cyclischen Verbindung in einem polaren Lösungsmittel wie Methanol mit Hilfe eines Metallhydrids, wie Natriumborhydrid, intervenierend in einer Menge entsprechend einem halb-reduzierendem Äquivalent; für die Herstellung der besagten Verbindungen der Formel (I), worin $X = \text{Ar(R)-Se-}$; oder


Erlauben der besagten, gemäss einem der obigen synthetischen Wege erhaltenen cyclischen Verbindung, sich bei Umgebungstemperatur mit der Thiol-Verbindung entsprechend den Werten von $X \neq \text{Ar(R)-Se-}$ umzusetzen, für die Herstellung der besagten Verbindungen der Formel (I), worin $X \neq \text{Ar(R)-Se-}$.

11. Verfahren zum Herstellen von Organoselenverbindungen gemäss Anspruch 1 oder 2 der allgemeinen Formel (I), worin $R = H$; also der allgemeinen Formel (II) unten:



dadurch gekennzeichnet, dass es die folgenden wesentlichen Schritte umfasst:

a/ Herstellen oder Verwenden eines Derivats der Formel (III):



N-Alkyliert, N-Aryliert, N-Acyliert oder N-Sulfonyliert wenn $Z = \text{NR}_5$; dann, gemäss der in Erwägung gezogenen Serien:

- für die Herstellung besagter Diselenidverbindungen der Formel (II), worin $R_8 = \text{Wasserstoff}$:

b1/ Erlauben, dass sich das besagte Derivat der Formel (III) mit einem elektrophilen Selenderivat in einem nicht-polaren Lösungsmittel umsetzt;

c1/ falls notwendig De-acylieren oder De-sulfonylieren der erhaltenen Verbindung gemäss üblicher Verfahren;

- für die Herstellung besagter Diselenidverbindungen der Formel (II), worin $R_8 = \text{nied.-Alkyl}$, gegebenenfalls substituiertes nied.-Alkyl, gegebenenfalls substituiertes nied.-Hetero-alkyl:

b2/ Alkylieren des besagten Derivats der Formel (III) in Position 2, indem man es zuerst mit einer Organolithiumbase wie Lithiumdiisopropylamid, dann mit einem Halid, insbesondere einem nied.-Alkyl, gegebenenfalls substituiertem nied.-Alkyl oder gegebenenfalls substituiertem nied.-Heteroalkyliodid behandelt;

c2/ Erlauben, dass sich das besagte Alkylderivat mit einem elektrophilen Selenderivat in einem nicht-

polaren Lösungsmittel umgesetzt;
d2/ falls notwendig, De-acylieren oder De-sulfonylieren der erhaltenen Verbindung gemäss üblicher Verfahren;

- 5 - für die Herstellung von besagten Diselenidverbindungen der Formel (II), in welchen R_8 = gegebenenfalls substituiertes Aryl, gegebenenfalls substituiertes Heteroaryl:

b3/ Behandeln des besagten Derivats der Formel (III) in einem nicht polaren, organischen Lösungsmittel in Anwesenheit einer starken Base mit einem Trialkylzinnhalid oder einem Zinkhalid, um das entsprechende Stannyl- oder Zinkderivat zu erhalten, dann Behandeln des besagten Stannyl- oder Zinkderivats mit einem halogenaromatischen Derivat in Anwesenheit von Palladium in einem nicht polaren organischen Lösungsmittel;

c3/ Erlauben, dass sich die erhaltene Verbindung mit einem elektrophilem Selenderivat in einem nicht-polaren Lösungsmittel umgesetzt;

15 d3/ falls notwendig De-acylieren oder De-sulfonylieren der erhaltenen Verbindung gemäss üblicher Verfahren;

b/ falls notwendig, Erlauben, dass sich die besagte, gemäss einem der obigen synthetischen Wege erhaltene Verbindung, in einem polaren Lösungsmittel mit einem passenden Mercaptan umgesetzt, um die entsprechende Selenosulfidverbindung zu erhalten.

Revendications

- 25 1. Composés d'organosélénium de formule générale (I) :



dans laquelle :

35

R = hydrogène ; $-C(R_1R_2)-A-B$;

40 R_1 = alkyle inférieur, linéaire ou ramifié, ayant de 1 à 8 atomes de carbone ; aryle éventuellement substitué ; aralkyle inférieur éventuellement substitué ; où le terme "aryle" signifie un groupe aromatique choisi parmi phényle et naphthyle ; où le terme "substitué" concernant les termes "aryle", "aralkyle" signifie que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C_1-C_8 , trifluorométhyle, alcoxy en C_1-C_8 , hydroxyle, nitro, amino, alkyle en C_1-C_8 -amino, dialkyle en C_1-C_8 -amino, sulfoxy, sulfonyl, sulfonamide, sulfoalkyle en C_1-C_8 , carboxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C_1-C_8 pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène, et où "aralkyle inférieur" signifie phényl-alkyle en C_1-C_8 ou naphthyl-alkyle en C_1-C_8 et hétéroaryl-alkyle en C_1-C_8 ;


50 R_2 = alkyle inférieur, linéaire ou ramifié, ayant de 1 à 8 atomes de carbone ; aryle éventuellement substitué ; aralkyle inférieur éventuellement substitué ; où le terme "aryle" signifie un groupe aromatique choisi parmi phényle et naphthyle ; où le terme "substitué" concernant les termes "aryle", "aralkyle" signifie que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C_1-C_8 , trifluorométhyle, alcoxy en C_1-C_8 , hydroxyle, nitro, amino, alkyle en C_1-C_8 -amino, dialkyle en C_1-C_8 -amino, sulfoxy, sulfonyl, sulfonamide, sulfoalkyle en C_1-C_8 , carboxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C_1-C_8 pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène ;

A = CO ; $(CR_3R_4)_n$;

55 B représente NR_5R_6 ; $N^+R_5R_6R_7Y^-$; OR_5 ; SR_5 ;

Ar = un groupe phényle éventuellement substitué ou un radical éventuellement substitué de formule :


5



10

où Z représente O ; S ; NR₅ ; quand R = -C(R₁R₂)-A-B
ou
Ar = un radical de formule

15




20

dans laquelle Z = O ; S ; NR₅ ; quand R est l'hydrogène ;
où "substitué" concernant les termes "phényle", "radical" (à 5 chaînons, incluant Z) signifie que les groupes
sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les
groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino, alkyle en C₁-C₈-amino,
dialkyle en C₁-C₈-amino, sulfoxy, sulfonyl, sulfonamide, sulfoalkyle en C₁-C₈, carboxyle, carbalcoxy, car-
bamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs
atomes d'halogène,
X = Ar(R)-Se- ; -S-glutathion ; -S-N-acétylcystéine ; -S-cystéine ; -S-pénicillamine ; -S-albumine ; -S-glucose ;

30

35

40



45

50

55

R₃ = hydrogène ; alkyle inférieur ; linéaire ou ramifié, ayant de 1 à 8 atomes de carbone ; aryle éventuellement
substitué ; aralkyle inférieur éventuellement substitué ; où le terme "aryle" signifie un groupe aromatique choisi
parmi phényle et naphthyle ; où le terme "substitué" concernant les termes "aryle", "aralkyle" signifie que les
groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou dif-
férents choisis parmi les groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino,
alkyle en C₁-C₈-amino, dialkyle en C₁-C₈-amino, sulfoxy, sulfonyl, sulfonamide, sulfoalkyle en C₁-C₈, car-
boxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou subs-
titués par un ou plusieurs atomes d'halogène, et où "aralkyle inférieur" signifie phényl-alkyle en C₁-C₈ ou
naphthyl-alkyle en C₁-C₈ et hétéroaryl-alkyle en C₁-C₈ ;
R₄ = hydrogène ; alkyle inférieur ; linéaire ou ramifié, ayant de 1 à 8 atomes de carbone ; aryle éventuellement
substitué ; aralkyle inférieur éventuellement substitué ; où le terme "aryle" signifie un groupe aromatique choisi

parmi phényle et naphthyle ; où le terme "substitué" concernant les termes "aryle", "aralkyle" signifie que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino, alkyle en C₁-C₈-amino, dialkyle en C₁-C₈-amino, sulfoxy, sulfonyle, sulfonamide, sulfoalkyle en C₁-C₈, carboxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène, et où "aralkyle inférieur" signifie phényl-alkyle en C₁-C₈ ou naphthyl-alkyle en C₁-C₈ et hétéroaryl-alkyle en C₁-C₈ ;


R₅ = hydrogène, alkyle inférieur, aryle éventuellement substitué ; aralkyle inférieur éventuellement substitué ; hétéroaryle éventuellement substitué ; hétéroaralkyle inférieur éventuellement substitué ; CO(alkyle inférieur) ; CO(aryle) ; SO₂(alkyle inférieur) ; SO₂(aryle) ; où "inférieur" dans "alkyle inférieur" ou "aralkyle inférieur", ou "hétéroaralkyle inférieur", ou dans "CO(alkyle inférieur)" ou dans "SO₂(alkyle inférieur)" signifie un groupe alkyle, linéaire ou ramifié, ayant de 1 à 8 atomes de carbone ; le terme "aryle" signifie un groupe aromatique choisi parmi les groupes phényle et naphthyle, le terme "hétéroaryle" signifie un groupe aromatique mono- ou bicyclique, chaque cycle, ou noyau, comprenant 5 ou 6 atomes et ledit cycle, ou noyau, ou les deux cycles, ou noyaux, comprenant dans leur squelette carboné de 1 à 3 hétéroatomes choisis parmi l'azote, l'oxygène et le soufre ; et le terme "substitué" concernant les termes "aryle", "aralkyle", "hétéroaryle", "hétéroaralkyle" signifie que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino, alkyle en C₁-C₈-amino, dialkyle en C₁-C₈-amino, sulfoxy, sulfonyle, sulfonamide, sulfoalkyle en C₁-C₈, carboxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène ; les termes "aralkyle inférieur" et "hétéroaralkyle inférieur" signifient phényl-alkyle en C₁-C₈ ou naphthyl-alkyle en C₁-C₈ et hétéroaryl-alkyle en C₁-C₈ respectivement ;

R₆ = hydrogène ; alkyle inférieur, linéaire ou ramifié ayant de 1 à 8 atomes de carbone ; aryle éventuellement substitué ; aralkyle inférieur éventuellement substitué ; hétéroaryle éventuellement substitué ; hétéroaralkyle inférieur éventuellement substitué ; le terme "aryle" signifie un groupe aromatique choisi parmi les groupes phényle et naphthyle, le terme "hétéroaryle" signifie un groupe aromatique mono- ou bicyclique, chaque cycle, ou noyau, comprenant 5 ou 6 atomes et ledit cycle, ou noyau, ou les deux cycles, ou noyaux, comprenant dans leur squelette carboné de 1 à 3 hétéroatomes choisis parmi l'azote, l'oxygène et le soufre ; le terme "substitué" concernant les termes "aryle", "aralkyle", "hétéroaryle", "hétéroaralkyle" signifient que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino, alkyle en C₁-C₈-amino, dialkyle en C₁-C₈-amino, sulfoxy, sulfonyle, sulfonamide, sulfoalkyle en C₁-C₈, carboxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène, les termes "aralkyle inférieur" et "hétéroaralkyle inférieur" signifient phényl-alkyle en C₁-C₈ ou naphthyl-alkyle en C₁-C₈ et hétéroaryl-alkyle en C₁-C₈ respectivement ;

R₇ = hydrogène ; alkyle inférieur, linéaire ou ramifié ayant de 1 à 8 atomes de carbone ; aryle éventuellement substitué ; aralkyle inférieur éventuellement substitué ; hétéroaryle éventuellement substitué ; hétéroaralkyle inférieur éventuellement substitué ; le terme "aryle" signifie un groupe aromatique choisi parmi les groupes phényle et naphthyle, le terme "hétéroaryle" signifie un groupe aromatique mono- ou bicyclique, chaque cycle, ou noyau, comprenant 5 ou 6 atomes et ledit cycle, ou noyau, ou les deux cycles, ou noyaux, comprenant dans leur squelette carboné de 1 à 3 hétéroatomes choisis parmi l'azote, l'oxygène et le soufre ; le terme "substitué" concernant les termes "aryle", "aralkyle", "hétéroaryle", "hétéroaralkyle" signifient que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino, alkyle en C₁-C₈-amino, dialkyle en C₁-C₈-amino, sulfoxy, sulfonyle, sulfonamide, sulfoalkyle en C₁-C₈, carboxyle, carbalcoxy, carbamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène, les termes "aralkyle inférieur" et "hétéroaralkyle inférieur" signifient phényl-alkyle en C₁-C₈ ou naphthyl-alkyle en C₁-C₈ et hétéroaryl-alkyle en C₁-C₈ respectivement ;

R₈ = hydrogène ; trifluorométhyle ; alkyle inférieur, linéaire ou ramifié ayant de 1 à 8 atomes de carbone ; aryle éventuellement substitué ; aralkyle inférieur éventuellement substitué ; hétéroaryle éventuellement substitué ; hétéroaralkyle inférieur éventuellement substitué ; le terme "aryle" signifie un groupe aromatique choisi parmi les groupes phényle et naphthyle, le terme "hétéroaryle" signifie un groupe aromatique mono- ou bicyclique, chaque cycle, ou noyau, comprenant 5 ou 6 atomes et ledit cycle, ou noyau, ou les deux cycles, ou noyaux, comprenant dans leur squelette carboné de 1 à 3 hétéroatomes choisis parmi l'azote, l'oxygène et le soufre ; le terme "substitué" concernant les termes "aryle", "aralkyle", "hétéroaryle", "hétéroaralkyle" signifient que les groupes en question sont substitués sur la partie aromatique par un ou plusieurs groupes identiques ou différents choisis parmi les groupes : alkyle en C₁-C₈, trifluorométhyle, alcoxy en C₁-C₈, hydroxyle, nitro, amino, alkyle

en C₁-C₈-amino, dialkyle en C₁-C₈-amino, sulfoyle, sulfonyle, sulfonamide, sulfoalkyle en C₁-C₈, carboxyle, carbalcoyle, carbamide (lesdits groupes alkyle en C₁-C₈ pouvant être linéaires ou ramifiés) ou substitués par un ou plusieurs atomes d'halogène, les termes "aralkyle inférieur" et "hétéroaralkyle inférieur" signifiant phényl-alkyle en C₁-C₈ ou naphtyl-alkyle en C₁-C₈ et hétéroaryl-alkyle en C₁-C₈ respectivement ;



n = 0 ou 1 ;

X⁺ représente le cation d'une base pharmaceutiquement acceptable ;

Y⁻ représente l'anion d'un acide pharmaceutiquement acceptable ;

et leurs sels d'acides ou de bases pharmaceutiquement acceptables ;

avec les conditions que :

quand

R = -C(R₁R₂)-(CR₃R₄)-B avec B = NR₅R₆ ou N⁺R₅R₆R₇Y⁻

et

X = Ar(R)-Se- avec Ar = phényle éventuellement substitué,

alors

-C(R₁R₂) est différent de (CR₃R₄) ;

et

quand

Ar = phényle et

R = -C(R₁R₂)-C(O)-B avec B = NH₂ ou NHCH₃ ou NHCH₂C₆H₅ ou NHC₆H₅

et

X = Ar(R)-Se-

alors R₁ et R₂ ne peuvent pas représenter simultanément un groupe méthyle.

2. Composés d'organosélénium de formule générale (I), selon la revendication 1, choisi dans le groupe consistant en :

le diséléniure de di[2-[2'-(1'-amino-2'-méthyl)propyl]phényle],

le dichlorhydrate de diséléniure de di[2-[2'-(1'-amino-2'-méthyl)propyl]phényle],

le diparatoluenesulfonate de diséléniure de di[2-[2'-(1'-ammonium-2'-méthyl)propyl]phényle],

le diséléniure de di[2-[2'-(1'-amino-2'-méthyl)propyl]-4-méthoxy]phényle,

le diséléniure de di[2-[2'-(1'-méthylamino-2'-méthyl)propyl]-phényle],

le dichlorhydrate de diséléniure de di[2-[2'-(1'-méthylamino-2'-méthyl)propyl]phényle],

- le diséléniure de di[2-[2'-(1'-diméthylamino-2'-méthyl)propyl]-phényle],
 le diparatoluènesulfonate de diséléniure de di[2-[2'-(1'-triméthylammonium-2'-méthyl)propyl]phényle],
 le diséléniure de di[3-[2'-(1'-amino-2'-méthyl)propyl]-2-thiényle],
 le diséléniure de di[2-[2'-(1'-amino-2'-méthyl)propyl]-3-thiényle],
 le séléniure de S-(N-acétyl-L-cystéinyl)-[2-[2'-(1'-amino-2'-méthyl)-propyl]phényle],
 le séléniure de S-glutathionyl-[2-[2'-(1'-amino-2'-méthyl)-propyl]-phényle],
 le bis-4(5)-[2-phényl-4(5)-séléno-1H-imidazole],
 le bis-4(5)-[[2-(4'-carbométhoxy)phényl]-4(5)-séléno-1H-imidazole],
 le bis-4(5)-[2-[(4'-carboxyl)phényl]-4(5)-séléno-1H-imidazole],
 le bis-4(5)-[2-[4'-carbo(2''-(4'''-méthyl-pipérazin-1''-yl)éthoxyphényl)]-4(5)séléno-1H-imidazole],
 le bis-5-[2-[(4'-trifluorométhyl)phényl]-5-séléno-1-méthyl-imidazole],
 le bis-5-[2-[(4'-carbométhoxy)phényl]-5-séléno-1-méthyl-imidazole],
 le bis-5-[2-[(4'-carboxyl)-phényl]-5-séléno-1-méthyl-imidazole].
3. Utilisation de composés d'organoséléniure, de formule générale (I), selon la revendication 1 ou 2, comme antioxydants, destinés en particulier à être ajoutés à des agents conservateurs de greffes pour la transplantation d'organes d'origine humaine ou animale comme le coeur, le foie, le rein et les poumons ; et en particulier pour la fabrication de compositions pharmaceutiques à activité antioxydante, destinées en particulier :
- * aux traitements de toute affection physiopathologique dans laquelle une surproduction d'hydroperoxydes cytotoxiques contribue aux dégradations fonctionnelles de cellules ou de tissus ; et comprenant plus particulièrement :
 - * le traitement des pathologies cardio- et cérébrovasculaires inflammatoires et/ou ischémiques, comme le traitement préventif et/ou curatif des resténoses artérielles consécutives à une angioplastie, le traitement préventif et/ou curatif de sténoses artérielles consécutives à des allogreffes d'artères, le traitement de la claudication intermittente chez des patients affectés d'ischémie obstructive des membres inférieurs, le traitement des accidents cérébro-vasculaires d'origine ischémique ;
 - * le traitement des pathologies digestives inflammatoires et/ou ischémiques, comme le traitement des inflammations aiguës de l'intestin (maladie de Crohn, rectocolite hémorragique) ;
 - * le traitement des pathologies respiratoires inflammatoires et/ou ischémiques, comme le traitement du syndrome de détresse respiratoire aigu de l'adulte (ARDS) et du syndrome de détresse respiratoire du nouveau-né (IRDS) ;
 - * le traitement des pathologies ophtalmiques inflammatoires et/ou ischémiques, comme le traitement du glaucome ;
 - * le traitement des cataractes ;
 - * le traitement des allergies ophtalmiques aiguës ;
 - * le traitement des dégradations de la rétine qui sont associées à une dégénérescence maculaire ;
 - * le traitement des infections virales provoquant une immuno-déficience, comme le traitement du SIDA ; et
 - * le traitement des fibroses consécutives à une radiothérapie.
4. Utilisation de composés d'organoséléniure, de formule générale (I), selon la revendication 1 ou 2, pour la fabrication de compositions pharmaceutiques destinées au traitement :
- * des maladies inflammatoires de l'intestin comme la maladie de Crohn ou la rectocolite hémorragique ;
 - * du syndrome de détresse respiratoire aigu de l'adulte et du syndrome de détresse respiratoire du nouveau-né ;
 - * des cataractes ;
 - * du SIDA ;
 - * des fibroses consécutives à une radiothérapie.
5. Composition pharmaceutique, ayant notamment une activité antioxydante, caractérisée en ce qu'elle comprend, comme ingrédient actif, au moins un composé d'organoséléniure, de formule générale (I), selon la revendication 1 ou 2, ou l'un de ses sels pharmaceutiquement acceptables d'un acide ou d'une base, incorporés éventuellement dans un excipient, support ou véhicule pharmaceutiquement acceptable.
6. Composition pharmaceutique selon la revendication 5, caractérisée en ce que ledit composé d'organoséléniure de formule générale (I) ci-dessus, éventuellement sous forme de sel, est présent en une quantité de 0,1 à 5 % en

masse, avantageusement de 0,1 à 1 % en masse, par rapport à la masse totale de la composition.

7. Composition pharmaceutique selon l'une des revendications 5 et 6, caractérisée en ce qu'elle est sous forme d'une dose unitaire comprenant de 1 à 500 mg d'au moins un composé d'organosélénium de formule générale (I).

8. Composition pharmaceutique selon l'une quelconque des revendications 5 à 7, caractérisée en ce qu'il s'agit d'une composition pharmaceutique pour le traitement de toute affection physiopathologique dans laquelle une surproduction d'hydroperoxydes cytotoxiques contribue aux dégradations fonctionnelles de cellules ou de tissus ; cette surproduction d'hydroperoxydes pouvant être due :

- à l'activation des voies métaboliques intracellulaires comme celles des flavine ou cytochrome P-450 oxygénases, celles des lipoxigénases et celles des monoamine oxydases ;
- à l'activation d'enzymes contenues dans des cellules endothéliales (xanthine oxydase, 15-lipoxigénase) ou dans les plaquettes sanguines (cyclooxygénase et 12-lipoxigénase) ;
- à l'activation, par des cytokines comme $\text{TNF-}\alpha$, de cellules inflammatoires et/ou immunitaires comme les neutrophiles, les macrophages ou les lymphocytes ;
- à une intoxication par un agent xénobiotique produisant des radicaux libres ;
- à une irradiation volontaire, par exemple pratiquée pendant une radiothérapie, ou une irradiation accidentelle.

9. Composition pharmaceutique selon l'une quelconque des revendications 5 à 6, caractérisée en ce qu'il s'agit d'une composition pharmaceutique pour le traitement de toute affection physiopathologique dans laquelle une surproduction d'hydroperoxydes cytotoxiques contribue aux dégradations fonctionnelles de cellules ou de tissus ; et plus particulièrement :

- * pour le traitement des pathologies cardio- et cérébro-vasculaires inflammatoires et/ou ischémiques, comme le traitement préventif et/ou curatif des resténoses artérielles consécutives à une angioplastie, le traitement préventif et/ou curatif de sténoses artérielles consécutives à des allogreffes d'artères, le traitement de la claudication intermittente chez des patients affectés d'ischémie obstructive des membres inférieurs, le traitement des accidents cérébro-vasculaires d'origine ischémique ;
- * pour le traitement des pathologies digestives inflammatoires et/ou ischémiques, comme le traitement des inflammations aiguës de l'intestin (maladie de Crohn, rectocolite hémorragique) ;
- * pour le traitement des pathologies respiratoires inflammatoires et/ou ischémiques, comme le traitement du syndrome de détresse respiratoire aigu de l'adulte (ARDS) et du syndrome de détresse respiratoire du nouveau-né (IRDS) ;
- * pour le traitement des pathologies ophtalmiques inflammatoires et/ou ischémiques, comme le traitement du glaucome ;
- * pour le traitement des cataractes ;
- * pour le traitement des allergies ophtalmiques aiguës ;
- * pour le traitement des dégradations de la rétine qui sont associées à une dégénérescence maculaire ;
- * pour le traitement des infections virales provoquant une immuno-déficience, comme le traitement du SIDA ; et
- * pour le traitement des fibroses consécutives à une radiothérapie.

10. Procédé de préparation de composés d'organosélénium selon la revendication 1 ou 2, de formule générale (I), où $\text{R} = -\text{C}(\text{R}_1\text{R}_2)-\text{A}-\text{B}$ ($\text{R} = \text{H}$) ; caractérisé en ce qu'il comprend les étapes essentielles suivantes :

a/ préparation ou utilisation d'un dérivé d'orthohalogéno(hétéro)arylacétonitrile, éventuellement mono- ou gem-disubstitué dans la position benzylique ; puis, selon la série considérée :

- pour la préparation desdits composés de formule (I) où $\text{A} = (\text{CR}_3\text{R}_4)_n$ avec $n = 0$ (A n'existe pas) :

b1/ hydrolyse dudit dérivé nitrile en un dérivé amide,

c1-1/ transformation de celui-ci en un dérivé amine par une réaction de transposition selon des procédés conventionnels,

d1-1/ réaction dudit dérivé amine avec un dérivé nucléophile du sélénium, produit éventuellement *in situ*, en présence d'un sel de cuivre Cu(I) dans un solvant organique polaire, pour conduire au dérivé (hétéro)arylisosélénazoline correspondant,

e1-1/ éventuellement, N-alkylation ou N-arylation ou N-acylation ou N-sulfonylation, selon des protocoles conventionnels, dudit dérivé (hétéro)arylisosélénazoline ;

- pour la préparation desdits composés de formule (I) où $A = CO$:

b1/ hydrolyse dudit dérivé nitrile en un dérivé amide,
 c1-2/ réaction dudit dérivé amide avec un dérivé nucléophile du sélénium, produit éventuellement *in situ*, en présence d'un sel de cuivre Cu(I), dans un solvant organique polaire, pour conduire au dérivé (hétéro)arylisosélénazone correspondant,
 d1-2/ éventuellement, N-alkylation ou N-arylation ou N-acylation ou N-sulfonylation, dudit dérivé (hétéro)arylisosélénazone ;

- pour la préparation desdits composés de formule (I) où $A = CH_2$:

b2/ réduction dudit dérivé nitrile en un dérivé amine, à l'aide de borane par exemple dans un solvant étheré comme le tétrahydrofurane par exemple,
 c2/ réaction dudit dérivé amine avec un dérivé nucléophile du sélénium, produit éventuellement *in situ*, en présence d'un sel de cuivre Cu(I), dans un solvant organique polaire, pour conduire au dérivé (hétéro)arylisosélénazine correspondant,
 d2/ éventuellement, N-alkylation ou N-arylation ou N-acylation ou N-sulfonylation, selon des protocoles conventionnels, dudit dérivé (hétéro)arylisosélénazine ;

- pour la préparation desdits composés de formule (I) où $A = (CR_3R_4) (\neq CH_2)$:


b3/ mise en oeuvre d'une mono- ou d'une bis-C-alkylation dudit dérivé nitrile selon des procédés conventionnels, à l'aide d'un dérivé d'organolithium par exemple dans un solvant étheré comme le tétrahydrofurane ;
 c3/ réaction du dérivé amine obtenu avec un dérivé nucléophile du sélénium, produit éventuellement *in situ*, en présence d'un sel de cuivre Cu(I), dans un solvant organique polaire, pour conduire au dérivé (hétéro)arylisosélénazine correspondant ;
 d3/ éventuellement N-alkylation ou N-arylation ou N-acylation ou N-sulfonylation, selon des protocoles conventionnels, dudit dérivé (hétéro)arylisosélénazine correspondant ;

b/ réduction, dans un solvant polaire comme le méthanol, du composé cyclique obtenu, selon l'une quelconque des voies de synthèse ci-dessus, à l'aide d'un hydrure métallique, comme le borohydrure de sodium, intervenant en une quantité correspondant à un demi équivalent réducteur ; pour la préparation desdits composés de formule (I) où $X = Ar(R)-Se-$;

ou


réaction dudit composé cyclique obtenu selon l'une quelconque des voies de synthèse ci-dessus, à la température ambiante, avec le composé thiol correspondant aux valeurs de $X \neq Ar(R)-Se-$, pour la préparation desdits composés de formule (I) où $X \neq Ar(R)-Se-$.

11. Procédé de préparation de composés d'organosélénium selon la revendication 1 ou 2, de formule générale (I) où $R = H$; c'est-à-dire de formule générale (II) ci-dessous :



caractérisé en ce qu'il comprend les étapes essentielles suivantes :

- a/ préparation ou utilisation d'un dérivé de formule (III) :



(III) ;

N-alkylé, N-arylé, N-acylé ou N-sulfonylé quand $Z = NR_5$; puis, selon la série considérée :

- pour la préparation desdits composés diséléniures de formule (II) où $R_8 = \text{hydrogène}$:

b1/ réaction dudit dérivé de formule (III) avec un dérivé électrophile du sélénium, dans un solvant non polaire ;

c1/ si nécessaire, désacylation ou désulfonylation, selon des protocoles courants, du composé obtenu ;

- pour la préparation desdits composés diséléniures de formule (II) où $R_8 = \text{alkyle inférieur, aralkyle inférieur éventuellement substitué, hétéroaralkyle inférieur éventuellement substitué}$:

b2/ alkylation dudit dérivé de formule (III) en position 2 par traitement tout d'abord avec une base d'organolithium comme le diisopropylamide de lithium, puis avec un halogénure, notamment un iodure d'alkyle inférieur, d'aralkyle inférieur éventuellement substitué ou d'hétéroaralkyle inférieur éventuellement substitué ;

c2/ réaction dudit dérivé alkyle avec un dérivé électrophile du sélénium, dans un solvant non polaire ;
d2/ si nécessaire, désacylation ou désulfonylation, selon des protocoles courants, du composé obtenu ;

- pour la préparation desdits composés diséléniures de formule (II) où $R_8 = \text{aryle éventuellement substitué, hétéroaryle éventuellement substitué}$:

b3/ traitement dudit dérivé de formule (III), dans un solvant organique non polaire, en présence d'une base forte, avec un halogénure de trialkylétain ou un halogénure de zinc pour obtenir le dérivé stannyle ou de zinc correspondant, puis traitement dudit dérivé stannyle ou de zinc avec un dérivé halogénoaromatique, en présence de palladium, dans un solvant organique non polaire ;

c3/ réaction du composé obtenu avec un dérivé électrophile du sélénium, dans un solvant non polaire ;
d3/ si nécessaire, désacylation ou désulfonylation, selon des protocoles courants, du composé obtenu ;

b/ si nécessaire, réaction dudit composé diséléniure obtenu selon l'une quelconque des voies de synthèse ci-dessus, dans un solvant polaire, avec un mercaptan adéquat ; pour obtenir le composé sélénosulfure correspondant.

Figure 1




Figure 2




Figure 3

